

● UNLICENSED USE

- ▶ In children *Fucibet*[®] *Lipid Cream* is not licensed for use in children under 6 years.

- **INTERACTIONS** → Appendix 1: corticosteroids · fusidate
- **PATIENT AND CARER ADVICE** Patients or carers should be counselled on application of betamethasone with fusidic acid preparations.

- **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.

Cream

CAUTIONARY AND ADVISORY LABELS 28

EXCIPIENTS: May contain Cetostearyl alcohol (including cetyl and stearyl alcohol), chlorocresol, hydroxybenzoates (parabens)

- ▶ **Fucibet** (LEO Pharma)

Betamethasone (as Betamethasone valerate) 1 mg per 1 gram, Fusidic acid 20 mg per 1 gram *Fucibet cream* | 30 gram [PoM] £6.38 DT = £6.38 | 60 gram [PoM] £12.76 DT = £12.76
Fucibet Lipid cream | 30 gram [PoM] £6.74 DT = £6.38

- ▶ **Xemacort** (Mylan)

Betamethasone (as Betamethasone valerate) 1 mg per 1 gram, Fusidic acid 20 mg per 1 gram *Xemacort 20mg/g / 1mg/g cream* | 30 gram [PoM] £6.05 DT = £6.38 | 60 gram [PoM] £12.45 DT = £12.76

Betamethasone with neomycin

The properties listed below are those particular to the combination only. For the properties of the components please consider, betamethasone p. 1309, neomycin sulfate p. 1296.

● INDICATIONS AND DOSE

Severe inflammatory skin disorders such as eczemas unresponsive to less potent corticosteroids | Psoriasis

- ▶ TO THE SKIN USING OINTMENT, OR TO THE SKIN USING CREAM
- ▶ Child 2–17 years: Apply 1–2 times a day, to be applied thinly
- ▶ Adult: Apply 1–2 times a day, to be applied thinly

POTENCY

- ▶ Betamethasone (as valerate) 0.1% with neomycin cream and ointment: potent.

- **INTERACTIONS** → Appendix 1: corticosteroids · neomycin
- **PATIENT AND CARER ADVICE** Patient counselling is advised for betamethasone with neomycin cream and ointment (application).

- **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.

Ointment

CAUTIONARY AND ADVISORY LABELS 28

- ▶ **Betamethasone with neomycin (Non-proprietary)**

Betamethasone (as Betamethasone valerate) 1 mg per 1 gram, Neomycin sulfate 5 mg per 1 gram *Betamethasone valerate 0.1% / Neomycin 0.5% ointment* | 30 gram [PoM] £38.88 DT = £31.36 | 100 gram [PoM] £97.00 DT = £104.52

Cream

CAUTIONARY AND ADVISORY LABELS 28

EXCIPIENTS: May contain Cetostearyl alcohol (including cetyl and stearyl alcohol), chlorocresol

- ▶ **Betamethasone with neomycin (Non-proprietary)**

Betamethasone (as Betamethasone valerate) 1 mg per 1 gram, Neomycin sulfate 5 mg per 1 gram *Betamethasone valerate 0.1% / Neomycin 0.5% cream* | 30 gram [PoM] £38.88 DT = £31.36 | 100 gram [PoM] £97.00 DT = £104.52

Betamethasone with salicylic acid

11-Mar-2020

The properties listed below are those particular to the combination only. For the properties of the components please consider, betamethasone p. 1309, salicylic acid p. 1353.

● INDICATIONS AND DOSE

DIPROSALIC[®] OINTMENT

Severe inflammatory skin disorders such as eczemas unresponsive to less potent corticosteroids | Psoriasis

- ▶ TO THE SKIN
- ▶ Child: Apply 1–2 times a day, max. 60 g per week
- ▶ Adult: Apply 1–2 times a day, max. 60 g per week

POTENCY

- ▶ Betamethasone (as dipropionate) 0.05% with salicylic acid 3%: potent.

DIPROSALIC[®] SCALP APPLICATION

Severe inflammatory skin disorders such as eczemas unresponsive to less potent corticosteroids | Psoriasis

- ▶ TO THE SKIN
- ▶ Child: Apply 1–2 times a day, apply a few drops
- ▶ Adult: Apply 1–2 times a day, apply a few drops

POTENCY

- ▶ Betamethasone (as dipropionate) 0.05% with salicylic acid 2%: potent.

- **INTERACTIONS** → Appendix 1: corticosteroids

- **PATIENT AND CARER ADVICE**

DIPROSALIC[®] OINTMENT Patients or carers should be counselled on application of betamethasone and salicylic acid preparations.

DIPROSALIC[®] SCALP APPLICATION Patients or carers should be counselled on application of betamethasone and salicylic acid scalp application.

- **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.

Ointment

CAUTIONARY AND ADVISORY LABELS 28

- ▶ **Diprosalic** (Merck Sharp & Dohme Ltd)

Betamethasone (as Betamethasone dipropionate) 500 microgram per 1 gram, Salicylic acid 30 mg per 1 gram *Diprosalic 0.05%/3% ointment* | 30 gram [PoM] £3.18 DT = £3.18 | 100 gram [PoM] £9.14 DT = £9.14

Liquid

CAUTIONARY AND ADVISORY LABELS 28

EXCIPIENTS: May contain Disodium edetate

- ▶ **Diprosalic** (Merck Sharp & Dohme Ltd)

Betamethasone (as Betamethasone dipropionate) 500 microgram per 1 ml, Salicylic acid 20 mg per 1 ml *Diprosalic 0.05%/2% scalp application* | 100 ml [PoM] £10.10 DT = £10.10

Clobetasol propionate with neomycin sulfate and nystatin

The properties listed below are those particular to the combination only. For the properties of the components please consider, clobetasol propionate p. 1310, neomycin sulfate p. 1296.

● INDICATIONS AND DOSE

Short-term treatment only of severe resistant inflammatory skin disorders such as recalcitrant eczemas associated with infection and unresponsive to less potent corticosteroids | Psoriasis associated with infection

- ▶ TO THE SKIN

- ▶ Adult: (consult product literature)

continued →