

● INDICATIONS AND DOSE

GLANDOSANE[®]

Carmellose sodium 500 mg, sorbitol 1.5 g, potassium chloride 60 mg, sodium chloride 42.2 mg, magnesium chloride 2.6 mg, calcium chloride 7.3 mg, and dipotassium hydrogen phosphate 17.1 mg/50 g, pH 5.75.

Dry mouth as a result of having (or having undergone) radiotherapy (ACBS) | Dry mouth as a result of sicca syndrome (ACBS)

- ▶ BY MOUTH
- ▶ Adult: Apply as required, spray onto oral and pharyngeal mucosa

● PROFESSION SPECIFIC INFORMATION

Dental practitioners' formulary

Glandosane[®] Aerosol Spray may be prescribed.

Glandosane synthetic saliva spray lemon (Fresenius Kabi Ltd)
50 ml • NHS indicative price = £5.68 • Drug Tariff (Part IXa)

Glandosane synthetic saliva spray natural (Fresenius Kabi Ltd)
50 ml • NHS indicative price = £5.68 • Drug Tariff (Part IXa)

Glandosane synthetic saliva spray peppermint (Fresenius Kabi Ltd)
50 ml • NHS indicative price = £5.68 • Drug Tariff (Part IXa)

● INDICATIONS AND DOSE

ORALIEVE MOISTURISING MOUTH SPRAY

Aqua, glycerin, xylitol, poloxamer 407, sodium benzoate, monosodium phosphate, xanthan gum, aroma, disodium phosphate, benzoic acid, whey protein, lactoferrin, lactoperoxidase, potassium thiocyanate, glucose oxidase, SLS free, alcohol free, pH 5.8

Dry mouth

- ▶ BY MOUTH
- ▶ Adult: 1 spray as required, on the inside of each cheek, gums and tongue

Oralieve moisturising mouth spray (Oralieve UK)
50 ml • NHS indicative price = £4.95 • Drug Tariff (Part IXa)

● INDICATIONS AND DOSE

ORALIEVE GEL

Symptomatic treatment of dry mouth

- ▶ BY MOUTH
- ▶ Adult: Apply as required, particularly at night, to oral mucosa

● PRESCRIBING AND DISPENSING INFORMATION

Contains traces of milk protein and egg white protein.

Oralieve moisturising mouth gel (Oralieve UK)
50 ml • NHS indicative price = £2.96 • Drug Tariff (Part IXa)

● INDICATIONS AND DOSE

SST[®]

Sugar-free, citric acid, malic acid and other ingredients in a sorbitol base.

Symptomatic treatment of dry mouth in patients with impaired salivary gland function and patent salivary ducts

- ▶ BY MOUTH
- ▶ Adult: 1 tablet as required, allow tablet to dissolve slowly in the mouth

● PROFESSION SPECIFIC INFORMATION

Dental practitioners' formulary

May be prescribed as Saliva Stimulating Tablets.

SST saliva stimulating tablets (Sinclair IS Pharma Plc)
100 tablet • NHS indicative price = £4.86 • Drug Tariff (Part IXa)

● INDICATIONS AND DOSE

SALIVEZE[®]

Carmellose sodium (sodium carboxymethylcellulose), calcium chloride, magnesium chloride, potassium chloride, sodium chloride, and dibasic sodium phosphate, pH neutral

Dry mouth as a result of having (or having undergone) radiotherapy (ACBS) | Dry mouth as a result of sicca syndrome (ACBS)

- ▶ BY MOUTH
- ▶ Adult: Apply 1 spray as required, spray onto oral mucosa

● PROFESSION SPECIFIC INFORMATION

Dental practitioners' formulary

Saliveze[®] Oral Spray may be prescribed.

Saliveze mouth spray (Wyvern Medical Ltd)
50 ml • NHS indicative price = £3.50 • Drug Tariff (Part IXa)

● INDICATIONS AND DOSE

SALIVIX[®]

Sugar-free, reddish-amber, acacia, malic acid and other ingredients.

Symptomatic treatment of dry mouth

- ▶ BY MOUTH USING PASTILLES
- ▶ Adult: 1 unit as required, suck pastille

● PROFESSION SPECIFIC INFORMATION

Dental practitioners' formulary

Salivix[®] Pastilles may be prescribed as Artificial Saliva Pastilles.

Salivix pastilles (Galen Ltd)
50 pastille • NHS indicative price = £3.59 • Drug Tariff (Part IXa)

● INDICATIONS AND DOSE

XEROTIN[®]

Sugar-free, water, sorbitol, carmellose (carboxymethylcellulose), potassium chloride, sodium chloride, potassium phosphate, magnesium chloride, calcium chloride and other ingredients, pH neutral.

Symptomatic treatment of dry mouth

- ▶ BY MOUTH
- ▶ Adult: 1 spray as required

● PROFESSION SPECIFIC INFORMATION

Dental practitioners' formulary

Xerotin[®] Oral Spray may be prescribed as Artificial Saliva Oral Spray.

Xerotin spray (SpePharm UK Ltd)
100 ml • NHS indicative price = £6.86 • Drug Tariff (Part IXa)

PARASYMPATHOMIMETICS

Pilocarpine

31-Oct-2019

● INDICATIONS AND DOSE

Xerostomia following irradiation for head and neck cancer

- ▶ BY MOUTH
- ▶ Adult: 5 mg 3 times a day for 4 weeks, then increased if tolerated to up to 30 mg daily in divided doses if required, dose to be taken with or immediately after meals (last dose always with evening meal), maximum therapeutic effect normally within 4–8 weeks; discontinue if no improvement after 2–3 months

Dry mouth and dry eyes in Sjögren's syndrome

- ▶ BY MOUTH
- ▶ Adult: 5 mg 4 times a day; increased if tolerated to up to 30 mg daily in divided doses if required, dose to be taken with meals and at bedtime, discontinue if no improvement after 2–3 months