

Liquid

EXCIPIENTS: May contain Cetostearyl alcohol (including cetyl and stearyl alcohol), hydroxybenzoates (parabens), propylene glycol

▶ **Dalacin T** (Pfizer Ltd)

Clindamycin (as Clindamycin phosphate) 10 mg per 1 ml Dalacin T 1% topical lotion | 30 ml [PoM] £5.08 DT = £5.08 | 60 ml [PoM] £10.16

Dalacin T 1% topical solution | 30 ml [PoM] £4.34 DT = £4.34 | 50 ml [PoM] £7.23

Combinations available: **Benzoyl peroxide with clindamycin**, below · **Tretinoin with clindamycin**, p. 1338

ANTIBACTERIALS > **MACROLIDES****Erythromycin with zinc acetate** 20-Apr-2020

The properties listed below are those particular to the combination only. For the properties of the components please consider, erythromycin p. 569.

● **INDICATIONS AND DOSE****Acne vulgaris**

- ▶ TO THE SKIN
- ▶ Child: Apply twice daily
- ▶ Adult: Apply twice daily

● **INTERACTIONS** → Appendix 1: macrolides● **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.**Liquid**▶ **Zineryt** (LEO Pharma)

Zinc acetate 12 mg per 1 ml, Erythromycin 40 mg per 1 ml Zineryt lotion | 30 ml [PoM] £9.25 DT = £9.25 | 90 ml [PoM] £20.02 DT = £20.02

ANTISEPTICS AND DISINFECTANTS > **PEROXIDES****Benzoyl peroxide**● **INDICATIONS AND DOSE****Acne vulgaris**

- ▶ TO THE SKIN
- ▶ Child 12-17 years: Apply 1–2 times a day, preferably apply after washing with soap and water, start treatment with lower-strength preparations
- ▶ Adult: Apply 1–2 times a day, preferably apply after washing with soap and water, start treatment with lower-strength preparations

● **CAUTIONS** Avoid contact with broken skin · avoid contact with eyes · avoid contact with mouth · avoid contact with mucous membranes · avoid excessive exposure to sunlight● **SIDE-EFFECTS**

- ▶ Common or very common Skin reactions
- ▶ Frequency not known Facial swelling

SIDE-EFFECTS, FURTHER INFORMATION Reduce frequency or suspend use until skin irritation subsides and re-introduce at reduced frequency.

● **PATIENT AND CARER ADVICE** May bleach fabrics and hair.● **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.**Gel**

EXCIPIENTS: May contain Fragrances, propylene glycol

▶ **Acneide** (Galderma (UK) Ltd)

Benzoyl peroxide 50 mg per 1 gram Acneide 5% gel | 30 gram [P] £5.44 DT = £5.44 | 60 gram [P] £10.68 DT = £10.68
Acneide Wash 5% gel | 50 gram [P] £5.44 DT = £5.44

Combinations available: **Adapalene with benzoyl peroxide**, p. 1337

Benzoyl peroxide with clindamycin

09-Mar-2020

The properties listed below are those particular to the combination only. For the properties of the components please consider, benzoyl peroxide above, clindamycin p. 1335.

● **INDICATIONS AND DOSE****Acne vulgaris**

- ▶ TO THE SKIN
- ▶ Child 12-17 years: Apply once daily, dose to be applied in the evening
- ▶ Adult: Apply once daily, dose to be applied in the evening

● **INTERACTIONS** → Appendix 1: clindamycin● **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.**Gel**

EXCIPIENTS: May contain Disodium edetate

▶ **Duac** (Stiefel Laboratories (UK) Ltd)

Clindamycin (as Clindamycin phosphate) 10 mg per 1 gram, Benzoyl peroxide 30 mg per 1 gram Duac Once Daily gel (3% and 1%) | 30 gram [PoM] £13.14 DT = £13.14 | 60 gram [PoM] £26.28
Clindamycin (as Clindamycin phosphate) 10 mg per 1 gram, Benzoyl peroxide 50 mg per 1 gram Duac Once Daily gel (5% and 1%) | 30 gram [PoM] £13.14 DT = £13.14 | 60 gram [PoM] £26.28 DT = £26.28

DERMATOLOGICAL DRUGS > **ANTICOMEDONALS****Azelaic acid**

10-Mar-2020

● **INDICATIONS AND DOSE****FINACEA[®]****Facial acne vulgaris**

- ▶ TO THE SKIN
- ▶ Child 12-17 years: Apply twice daily, discontinue if no improvement after 1 month
- ▶ Adult: Apply twice daily, discontinue if no improvement after 1 month

Papulopustular rosacea

- ▶ TO THE SKIN
- ▶ Adult: Apply twice daily, discontinue if no improvement after 2 months

SKINOREN[®]**Acne vulgaris**

- ▶ TO THE SKIN
- ▶ Child 12-17 years: Apply twice daily
- ▶ Adult: Apply twice daily

Acne vulgaris in patients with sensitive skin

- ▶ TO THE SKIN
- ▶ Child 12-17 years: Apply once daily for 1 week, then apply twice daily
- ▶ Adult: Apply once daily for 1 week, then apply twice daily

● **CAUTIONS** Avoid contact with eyes · avoid contact with mouth · avoid contact with mucous membranes● **SIDE-EFFECTS**

- ▶ Uncommon Skin reactions
- ▶ Rare or very rare Asthma exacerbated · cheilitis
- ▶ Frequency not known Angioedema · eye swelling

● **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.**Cream**

EXCIPIENTS: May contain Propylene glycol

▶ **Skinoren** (LEO Pharma)

Azelaic acid 200 mg per 1 gram Skinoren 20% cream | 30 gram [PoM] £4.49 DT = £4.49