

Hypertensive emergencies

Hypertensive emergencies in children may be accompanied by signs of hypertensive encephalopathy, including seizures. Controlled reduction in blood pressure over 72–96 hours is essential; rapid reduction can reduce perfusion leading to organ damage. Treatment should be initiated with intravenous drugs; once blood pressure is controlled, oral therapy can be started. It may be necessary to infuse fluids particularly during the first 12 hours to expand plasma volume should the blood pressure drop too rapidly.

Controlled reduction of blood pressure is achieved by intravenous administration of labetalol hydrochloride or sodium nitroprusside p. 120. Esmolol hydrochloride p. 108 is useful for short-term use and has a short duration of action. Nicardipine hydrochloride p. 110 can be administered as a continuous intravenous infusion for life-threatening hypertension in paediatric intensive care settings. In less severe cases, nifedipine capsules can be used.

Other antihypertensive drugs which can be given intravenously include hydralazine hydrochloride and clonidine hydrochloride.

Hypertension in acute nephritis occurs as a result of sodium and water retention; it should be treated with sodium and fluid restriction, and with furosemide p. 142; antihypertensive drugs may be added if necessary.

Also see advice on short-term management of hypertensive episodes in phaeochromocytoma.

Phaeochromocytoma

Long-term management of phaeochromocytoma involves surgery. However, surgery should not take place until there is adequate blockade of both alpha- and beta-adrenoceptors. Alpha-blockers are used in the short-term management of hypertensive episodes in phaeochromocytoma. Once alpha blockade is established, tachycardia can be controlled by the cautious addition of a beta-blocker; a cardioselective beta-blocker is preferred. There is no nationwide consensus on the optimal drug regimen or doses used for the management of phaeochromocytoma.

Phenoxybenzamine hydrochloride p. 119, a powerful alpha-blocker, is effective in the management of phaeochromocytoma but it has many side-effects.

Pulmonary hypertension

Only pulmonary *arterial* hypertension is currently suitable for drug treatment. Pulmonary arterial hypertension includes persistent pulmonary hypertension of the newborn, idiopathic pulmonary arterial hypertension in children, and pulmonary hypertension related to congenital heart disease and cardiac surgery.

Some types of pulmonary hypertension are treated with vasodilator antihypertensive therapy and oxygen. Diuretics may also have a role in children with right-sided heart failure.

Initial treatment of *persistent pulmonary hypertension of the newborn* involves the administration of **nitric oxide**; epoprostenol p. 120 can be used until nitric oxide is available. Oral sildenafil p. 122 may be helpful in less severe cases. Epoprostenol and sildenafil can cause profound systemic hypotension. In rare circumstances either tolazoline p. 123 or magnesium sulfate p. 618 can be given by intravenous infusion when nitric oxide and epoprostenol have failed.

Treatment of *idiopathic pulmonary arterial hypertension* is determined by acute vasodilator testing; drugs used for treatment include calcium-channel blockers (usually nifedipine), long-term intravenous epoprostenol, nebulised iloprost p. 121, bosentan p. 121, or sildenafil. Anticoagulation (usually with warfarin sodium p. 99) may also be required to prevent secondary thrombosis.

Inhaled nitric oxide is a potent and selective pulmonary vasodilator. It acts on cyclic guanosine monophosphate

(cGMP) resulting in smooth muscle relaxation. Inhaled nitric oxide is used in the treatment of persistent pulmonary hypertension of the newborn, and may also be useful in other forms of arterial pulmonary hypertension. Dependency can occur with high doses and prolonged use; to avoid rebound pulmonary hypertension the drug should be withdrawn gradually, often with the aid of sildenafil p. 122.

Excess nitric oxide can cause methaemoglobinaemia; therefore, methaemoglobin concentration should be measured regularly, particularly in neonates.

Nitric oxide increases the risk of haemorrhage by inhibiting platelet aggregation, but it does not usually cause bleeding.

Epoprostenol p. 120 (prostacyclin) is a prostaglandin and a potent vasodilator. It is used in the treatment of persistent pulmonary hypertension of the newborn, idiopathic pulmonary arterial hypertension, and in the acute phase following cardiac surgery. It is given by continuous 24-hour intravenous infusion.

Epoprostenol is a powerful inhibitor of platelet aggregation and there is a possible risk of haemorrhage. It is sometimes used as an antiplatelet in renal dialysis when heparins are unsuitable or contra-indicated. It can also cause serious systemic hypotension and, if withdrawn suddenly, can cause pulmonary hypertensive crisis.

Children on prolonged treatment can become tolerant to epoprostenol, and therefore require an increase in dose.

Iloprost p. 121 is a synthetic analogue of epoprostenol and is efficacious when nebulised in adults with pulmonary arterial hypertension, but experience in children is limited. It is more stable than epoprostenol and has a longer half-life.

Bosentan p. 121 is a dual endothelin receptor antagonist used orally in the treatment of pulmonary arterial hypertension. The concentration of endothelin, a potent vasoconstrictor, is raised in sustained pulmonary hypertension.

Sildenafil, a vasodilator developed for the treatment of erectile dysfunction, is also used for pulmonary arterial hypertension. It is used either alone or as an adjunct to other drugs.

Sildenafil is a selective phosphodiesterase type-5 inhibitor. Inhibition of this enzyme in the lungs enhances the vasodilatory effects of nitric oxide and promotes relaxation of vascular smooth muscle.

Sildenafil has also been used in pulmonary hypertension for weaning children off inhaled nitric oxide following cardiac surgery, and less successfully in idiopathic pulmonary arterial hypertension.

Tolazoline p. 123 is now rarely used to correct pulmonary artery vasospasm in pulmonary hypertension of the newborn as better alternatives are available. Tolazoline is an alpha-blocker and produces both pulmonary and systemic vasodilation.

Advanced Pharmacy Services

Children with hypertension may be eligible for the New Medicines Service / Medicines Use Review service provided by a community pharmacist. For further information, see *Advanced Pharmacy Services* in Guidance on prescribing p. 1.

Antihypertensive drugs

Vasodilator antihypertensive drugs

Vasodilators have a potent hypotensive effect, especially when used in combination with a beta-blocker and a thiazide. **Important:** see Hypertension (hypertensive emergencies) for a warning on the hazards of a very rapid fall in blood pressure.

Hydralazine hydrochloride p. 118 is given by mouth as an adjunct to other antihypertensives for the treatment of