

Meningococcal vaccine

Overview

Almost all childhood meningococcal disease in the UK is caused by *Neisseria meningitidis* serogroups B and C.

Meningococcal group C conjugate vaccine protects only against infection by serogroup C and **meningococcal group B vaccine** protects only against infection by serogroup B.

The risk of meningococcal disease declines with age—immunisation is not generally recommended after the age of 25 years.

Tetravalent meningococcal vaccines that cover serogroups A, C, W135, and Y are available. Although the duration of protection has not been established, the **meningococcal groups A, C, W135, and Y conjugate vaccine** is likely to provide longer-lasting protection than the unconjugated meningococcal polysaccharide vaccine. The antibody response to serogroup C in unconjugated meningococcal polysaccharide vaccines in young children may be suboptimal [not currently available in the UK].

Meningococcal group B vaccines, *Bexsero*[®] and *Trumenba*[®], are licensed in the UK against infection caused by *Neisseria meningitidis* serogroup B. The use of *Bexsero*[®] is recommended in the Immunisation Schedule. *Bexsero*[®] contains 3 recombinant *Neisseria meningitidis* serogroup B proteins and the outer membrane vesicles from the NZ 98/254 strain, in order to achieve broad protection against *Neisseria meningitidis* serogroup B. *Trumenba*[®] contains 2 recombinant *Neisseria meningitidis* serogroup B proteins. The proteins are adsorbed onto an aluminium compound to stimulate an enhanced immune response.

Childhood immunisation

Meningococcal group C conjugate vaccine provides long-term protection against infection by serogroup C of *Neisseria meningitidis*. Immunisation consists of 1 dose given at 12 months of age (as the haemophilus influenzae type b with meningococcal group C vaccine p. 818) and a second dose given at 13–15 years of age (as the meningococcal groups A with C and W135 and Y vaccine p. 819) (see Immunisation Schedule).

Meningococcal group B vaccine provides protection against infection by serogroup B of *Neisseria meningitidis*. Immunisation consists of 1 dose given at 2 months of age, a second dose at 4 months of age, and a booster dose at 12 months of age (see *Immunisation Schedule* above).

Unimmunised children aged under 12 months should be given 1 dose of meningococcal group B vaccine (rDNA, component, adsorbed) p. 818 followed by a second dose two months later. They should then be vaccinated according to the Immunisation Schedule (ensuring at least a two month interval between doses of meningococcal group B vaccines). Unimmunised children aged 12–23 months should be given 2 doses of meningococcal group B vaccine (rDNA, component, adsorbed) separated by an interval of two months if they have received less than 2 doses in the first year of life. Unimmunised children aged 2–9 years should be given a single dose of meningococcal group C vaccine (as the haemophilus influenzae type b with meningococcal group C vaccine), followed by a booster dose of meningococcal groups A with C and W135 and Y vaccine at 13–15 years of age.

From 2015, unimmunised individuals aged 10–25 years, including those aged under 25 years who are attending university for the first time, should be given a single dose of meningococcal groups A with C and W135 and Y vaccine; a booster dose is not required.

Children with confirmed serogroup C disease, who have previously been immunised with meningococcal group C vaccine, should be offered meningococcal group C conjugate vaccine before discharge from hospital.

Travel

Individuals travelling to countries of risk should be immunised with meningococcal groups A, C, W135, and Y conjugate vaccine, even if they have previously received meningococcal group C conjugate vaccine. If an individual has recently received meningococcal group C conjugate vaccine, an interval of at least 4 weeks should be allowed before administration of the tetravalent (meningococcal groups A, C, W135, and Y) vaccine.

Vaccination is particularly important for those living or working with local people or visiting an area of risk during outbreaks.

Immunisation recommendations and requirements for visa entry for individual countries should be checked before travelling, particularly to countries in Sub-Saharan Africa, Asia, and the Indian sub-continent where epidemics of meningococcal outbreaks and infection are reported. Country-by-country information is available from the National Travel Health Network and Centre (www.nathnac.org/).

Proof of vaccination with the tetravalent (meningococcal groups A, C, W135, and Y) vaccine is required for those travelling to Saudi Arabia during the Hajj and Umrah pilgrimages (where outbreaks of the W135 strain have occurred).

Contacts

For advice on the immunisation of *laboratory workers and close contacts* of cases of meningococcal disease in the UK and on the role of the vaccine in the control of *local outbreaks*, consult Guidelines for Public Health Management of Meningococcal Disease in the UK at www.gov.uk/phe. Also see antibacterial prophylaxis for prevention of secondary cases of meningococcal meningitis.

Useful Resources

Advice reflects that in the handbook Immunisation against Infectious Disease (2013), which in turn reflects the guidance of the Joint Committee on Vaccination and Immunisation (JCVI). The advice also incorporates changes announced by the Chief Medical Officer and Health Department Updates. Chapters from the handbook (including updates since 2013) are available at:

www.gov.uk/government/collections/immunisation-against-infectious-disease-the-green-book

Pertussis vaccine

Overview

Pertussis vaccine is given as a combination preparation containing other vaccines. Acellular vaccines are derived from highly purified components of *Bordetella pertussis*. Primary immunisation against pertussis (whooping cough) requires 3 doses of an acellular pertussis-containing vaccine (see Immunisation schedule), given at intervals of 1 month from the age of 2 months.

All children up to the age of 10 years should receive primary immunisation with a combination vaccine of diphtheria with tetanus, pertussis, hepatitis B, poliomyelitis and haemophilus influenzae type b vaccine p. 816 (*Infanrix hexa*[®]).

A booster dose of an acellular pertussis-containing vaccine should ideally be given 3 years after the primary course, although, the interval can be reduced to 1 year if the primary course was delayed.

Children aged 1–10 years who have not received a *pertussis-containing* vaccine as part of their primary immunisation should be offered 1 dose of a suitable pertussis-containing vaccine; after an interval of at least 1 year, a booster dose of a suitable pertussis-containing vaccine should be given. Immunisation against pertussis is