

- **Zeroneum** (Thornton & Ross Ltd)
Soya oil 833.5 mg per 1 gram Zeroneum 83.35% bath additive | 500 ml £4.48

Emollient bath and shower products, tar-containing

17-Aug-2016

● INDICATIONS AND DOSE

POLYTAR EMOLLIENT[®]

Psoriasis, eczema, atopic and pruritic dermatoses

▶ TO THE SKIN

- Child: 2–4 capfuls/bath, add 15–30 mL to an adult-size bath and proportionally less for a child's bath; soak for 20 minutes

PSORIDERM[®] EMULSION

Psoriasis

▶ TO THE SKIN

- Child: Up to 30 mL/bath, use 30mL in adult-size bath, and proportionately less for a child's bath, soak for 5 minutes

IMPORTANT SAFETY INFORMATION

These preparations make skin and surfaces slippery—particular care is needed when bathing.

MHRA/CHM ADVICE (UPDATED DECEMBER 2018): EMOLLIENTS: NEW INFORMATION ABOUT RISK OF SEVERE AND FATAL BURNS WITH PARAFFIN-CONTAINING AND PARAFFIN-FREE EMOLLIENTS See Emollient and barrier preparations p. 737.

- **DIRECTIONS FOR ADMINISTRATION** Emollient bath additives should be added to bath water; hydration can be improved by soaking in the bath for 10–20 minutes. Some bath emollients can be applied to wet skin undiluted and rinsed off. Emollient preparations contained in tubs should be removed with a clean spoon or spatula to reduce bacterial contamination of the emollient. Emollients should be applied in the direction of hair growth to reduce the risk of folliculitis.

- **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.

Bath additive

CAUTIONARY AND ADVISORY LABELS 15

EXCIPIENTS: May contain Isopropyl palmitate, polysorbates

- **Psoriderm** (Dermal Laboratories Ltd)
Coal tar distilled 400 mg per 1 ml Psoriderm Emulsion 40% bath additive | 200 ml [P] £2.74 DT = £2.74

Emollient creams and ointments, antimicrobial-containing

17-Aug-2016

● INDICATIONS AND DOSE

Dry and pruritic skin conditions including eczema and dermatitis

▶ TO THE SKIN

- Child: To be applied to the skin or used as a soap substitute

IMPORTANT SAFETY INFORMATION

These preparations make skin and surfaces slippery—particular care is needed when bathing.

MHRA/CHM ADVICE (UPDATED DECEMBER 2018): EMOLLIENTS: NEW INFORMATION ABOUT RISK OF SEVERE AND FATAL BURNS WITH PARAFFIN-CONTAINING AND PARAFFIN-FREE EMOLLIENTS See Emollient and barrier preparations p. 737.

- **DIRECTIONS FOR ADMINISTRATION** Emollients should be applied immediately after washing or bathing to maximise

the effect of skin hydration. Emollient preparations contained in tubs should be removed with a clean spoon or spatula to reduce bacterial contamination of the emollient. Emollients should be applied in the direction of hair growth to reduce the risk of folliculitis.

● PRESCRIBING AND DISPENSING INFORMATION

Preparations containing an antibacterial should be avoided unless infection is present or is a frequent complication.

- **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.

Cream

CAUTIONARY AND ADVISORY LABELS 15

EXCIPIENTS: May contain Cetostearyl alcohol (including cetyl and stearyl alcohol)

- **Dermol** (Dermal Laboratories Ltd)

Benzalkonium chloride 1 mg per 1 gram, Chlorhexidine hydrochloride 1 mg per 1 gram, Isopropyl myristate 100 mg per 1 gram, Liquid paraffin 100 mg per 1 gram Dermol cream | 100 gram [P] £2.86 | 500 gram [P] £6.63

Liquid

CAUTIONARY AND ADVISORY LABELS 15

EXCIPIENTS: May contain Cetostearyl alcohol (including cetyl and stearyl alcohol)

- **Dermol 500** (Dermal Laboratories Ltd)

Benzalkonium chloride 1 mg per 1 gram, Chlorhexidine hydrochloride 1 mg per 1 gram, Isopropyl myristate 25 mg per 1 gram, Liquid paraffin 25 mg per 1 gram Dermol 500 lotion | 500 ml [P] £6.04

Emollient creams and ointments, colloidal oatmeal-containing

15-Jan-2019

● INDICATIONS AND DOSE

Endogenous and exogenous eczema | Xeroderma | Ichthyosis

▶ TO THE SKIN

- Child: (consult product literature)

IMPORTANT SAFETY INFORMATION

MHRA/CHM ADVICE (UPDATED DECEMBER 2018): EMOLLIENTS: NEW INFORMATION ABOUT RISK OF SEVERE AND FATAL BURNS WITH PARAFFIN-CONTAINING AND PARAFFIN-FREE EMOLLIENTS See Emollient and barrier preparations p. 737.

- **DIRECTIONS FOR ADMINISTRATION** Emollients should be applied immediately after washing or bathing to maximise the effect of skin hydration. Emollient preparations contained in tubs should be removed with a clean spoon or spatula to reduce bacterial contamination of the emollient. Emollients should be applied in the direction of hair growth to reduce the risk of folliculitis.

- **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.

Cream and lotion

CAUTIONARY AND ADVISORY LABELS 15

EXCIPIENTS: May contain Benzyl alcohol, cetostearyl alcohol (including cetyl and stearyl alcohol), isopropyl palmitate

- **Aveeno** (Johnson & Johnson Ltd)

Aveeno lotion | 500 ml (ACBS) £6.66

Aveeno cream | 100 ml (ACBS) £3.97 | 300 ml (ACBS) £6.80 | 500 ml (ACBS) £6.47