

alternate days for a total of 3 doses provides protection 7–10 days after the last dose. If travelling from a non-endemic area to an area where typhoid is endemic, a booster consisting of 3 doses is recommended every 3 years. The oral typhoid vaccine should be avoided in immunosuppressed and HIV-infected children.

Prevention of typhoid primarily depends on improving sanitation and water supplies in endemic areas and on scrupulous personal, food and water hygiene.

All suspected cases of typhoid fever must be notified to the local health protection unit. Where there is a community level outbreak, specialist advice should be sought from Public Health England (tel. 020 8200 4400) or, in Scotland, Health Protection Scotland (tel. 0140 300 1191).

### Useful Resources

Recommendations reflect Chapter 33, Typhoid, in *Immunisation against infectious disease*—‘The Green Book’. Public Health England, August 2015.

[www.gov.uk/government/publications/typhoid-the-green-book-chapter-33](http://www.gov.uk/government/publications/typhoid-the-green-book-chapter-33)

National Travel Health Network and Centre  
[nathnac.net](http://nathnac.net)

## Varicella-zoster vaccine

### Overview

Varicella-zoster vaccine p. 828 (live) is licensed for immunisation against varicella (chickenpox) in seronegative individuals. It is not recommended for routine use in children but can be given to seronegative healthy children over 1 year who come into close contact with individuals at high risk of severe varicella infections.

Rarely, the varicella-zoster vaccine virus has been transmitted from the vaccinated individual to close contacts. Therefore, contact with the following should be avoided if a vaccine-related cutaneous rash develops within 4–6 weeks of the first or second dose:

- varicella-susceptible pregnant females;
- individuals at high risk of severe varicella, including those with immunodeficiency or those receiving immunosuppressive therapy.

Varicella-zoster immunoglobulin p. 798 is used to protect susceptible children at increased risk of severe varicella infection (see Immunoglobulins p. 793).

### Useful Resources

Advice reflects that in the handbook *Immunisation against Infectious Disease* (2013), which in turn reflects the guidance of the Joint Committee on Vaccination and Immunisation (JCVI). The advice also incorporates changes announced by the Chief Medical Officer and Health Department Updates. Chapters from the handbook (including updates since 2013) are available at:

[www.gov.uk/government/collections/immunisation-against-infectious-disease-the-green-book](http://www.gov.uk/government/collections/immunisation-against-infectious-disease-the-green-book)

## Yellow fever vaccine

19-Mar-2019

### Overview

Yellow fever vaccine, live p. 829 is an attenuated preparation of yellow fever virus grown in chick eggs. Yellow fever vaccine, live is recommended for:

- children aged 9 months or older who are travelling to, or living in areas or countries with a risk of yellow fever transmission;
- children aged 9 months or older who are travelling to, or living in countries that require an International Certificate

of Vaccination or Prophylaxis (ICVP) for entry (information about countries at risk of yellow fever is available from the National Health Network and Centre).

Children aged under 9 months are at risk of vaccine-associated encephalitis, with the risk being inversely proportional to age. Children aged under 6 months should **not** be vaccinated. Children aged 6–9 months should only be vaccinated following a detailed risk assessment, and vaccination is generally only recommended if the risk of yellow fever transmission is high (such as during epidemics/outbreaks). If travel is unavoidable, seek expert advice on whether to vaccinate.

A single-dose of yellow fever vaccine, live confers life-long immunity against yellow fever disease. Immunisation should be performed at least 10 days before travelling to an endemic area to allow protective immunity to develop and for the ICVP (if required) to become valid.

Reinforcing immunisation is not needed, except for a small subset of children at continued risk who may not have developed long-term protection from their initial yellow fever vaccine, live vaccination—seek expert advice.

Yellow fever vaccine, live should be avoided in HIV-infected and immunosuppressed children. If the yellow fever risk is unavoidable, consult the Children’s HIV Association of UK and Ireland ([www.chiva.org.uk/guidelines/immunisation](http://www.chiva.org.uk/guidelines/immunisation)) or other specialist advice.

All suspected cases of yellow fever must be notified to the local health protection unit. Where there is a community level outbreak, specialist advice should be sought from Public Health England (tel. 020 8200 4400) or, in Scotland, Health Protection Scotland (tel. 0140 300 1191).

### Useful Resources

Recommendations reflect Chapter 35, Yellow fever, in *Immunisation against infectious disease*—‘The Green book’. Public Health England, January 2019.

[www.gov.uk/government/publications/yellow-fever-the-green-book-chapter-35](http://www.gov.uk/government/publications/yellow-fever-the-green-book-chapter-35)

National Travel Health Network and Centre  
[nathnac.net](http://nathnac.net)

## Vaccines for travel

### Immunisation for travel

See advice on Malaria, treatment p. 407.

No special immunisation is required for travellers to the United States, Europe, Australia, or New Zealand, although all travellers should have immunity to tetanus and poliomyelitis (and childhood immunisations should be up to date); Tick-borne encephalitis vaccine is recommended for immunisation of those working in, or visiting, high-risk areas. Certain special precautions are required in non-European areas surrounding the Mediterranean, in Africa, the Middle East, Asia, and South America.

Travellers to areas that have a high incidence of **poliomyelitis** or **tuberculosis** should be immunised with the appropriate vaccine; in the case of poliomyelitis previously immunised travellers may be given a booster dose of a preparation containing inactivated poliomyelitis vaccine. BCG immunisation is recommended for travellers aged under 16 years proposing to stay for longer than 3 months (or in close contact with the local population) in countries with an incidence of tuberculosis greater than 40 per 100 000 (list of countries where the incidence of tuberculosis is greater than 40 cases per 100 000 is available from [www.gov.uk/phen](http://www.gov.uk/phen)); it should preferably be given 3 months or more before departure.

**Yellow fever** immunisation is recommended for travel to the endemic zones of Africa and South America. Many countries require an International Certificate of Vaccination