

Cinchocaine with hydrocortisone

21-Dec-2017

● INDICATIONS AND DOSE

PROCTOSEDL[®] OINTMENT

Haemorrhoids | Pruritus ani

- ▶ TO THE SKIN, OR BY RECTUM
- ▶ Child: Apply twice daily, to be administered morning and night and after a bowel movement. Apply externally or by rectum. Do not use for longer than 7 days

PROCTOSEDL[®] SUPPOSITORIES

Haemorrhoids | Pruritus ani

- ▶ BY RECTUM
- ▶ Child 12–17 years: 1 suppository, insert suppository night and morning and after a bowel movement. Do not use for longer than 7 days

UNIROID-HC[®] OINTMENT

Haemorrhoids | Pruritus ani

- ▶ TO THE SKIN, OR BY RECTUM
- ▶ Child 1 month–11 years (under medical advice only): Apply twice daily, and apply after a bowel movement, apply externally or by rectum, do not use for longer than 7 days
- ▶ Child 12–17 years: Apply twice daily, and apply after a bowel movement, apply externally or by rectum, do not use for longer than 7 days

UNIROID-HC[®] SUPPOSITORIES

Haemorrhoids | Pruritus ani

- ▶ BY RECTUM
- ▶ Child 12–17 years: 1 suppository, insert twice daily and after a bowel movement. Do not use for longer than 7 days

IMPORTANT SAFETY INFORMATION

MHRA/CHM ADVICE: CORTICOSTEROIDS: RARE RISK OF CENTRAL SEROUS CHORIOPATHY WITH LOCAL AS WELL AS SYSTEMIC ADMINISTRATION (AUGUST 2017)

See Corticosteroids, general use p. 450.

- **CAUTIONS** Local anaesthetic component can be absorbed through the rectal mucosa (avoid excessive application, particularly in children and infants) · local anaesthetic component may cause sensitisation (use for short periods only—no longer than a few days)

- **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.

Ointment

- ▶ **Proctosedyl** (Sanofi)
Cinchocaine hydrochloride 5 mg per 1 gram, Hydrocortisone 5 mg per 1 gram Proctosedyl ointment | 30 gram [PoM] £10.34 DT = £10.34

- ▶ **Uniroid HC** (Chemidex Pharma Ltd)
Cinchocaine hydrochloride 5 mg per 1 gram, Hydrocortisone 5 mg per 1 gram Uniroid HC ointment | 30 gram [PoM] £4.23 DT = £10.34

Suppository

- ▶ **Proctosedyl** (Sanofi)
Cinchocaine hydrochloride 5 mg, Hydrocortisone 5 mg Proctosedyl suppositories | 12 suppository [PoM] £5.08 DT = £5.08
- ▶ **Uniroid HC** (Chemidex Pharma Ltd)
Cinchocaine hydrochloride 5 mg, Hydrocortisone 5 mg Uniroid HC suppositories | 12 suppository [PoM] £1.91 DT = £5.08

Cinchocaine with prednisolone

21-Dec-2017

● INDICATIONS AND DOSE

Haemorrhoids | Pruritus ani

- ▶ BY RECTUM USING OINTMENT
- ▶ Child: Apply twice daily for 5–7 days, apply 3–4 times a day on the first day if necessary, then apply once daily for a few days after symptoms have cleared
- ▶ BY RECTUM USING SUPPOSITORIES
- ▶ Child 12–17 years: 1 suppository daily for 5–7 days, to be inserted after a bowel movement

Haemorrhoids (severe cases) | Pruritus ani (severe cases)

- ▶ BY RECTUM USING SUPPOSITORIES
- ▶ Child 12–17 years: Initially 1 suppository 2–3 times a day, then 1 suppository daily for a total of 5–7 days, to be inserted after a bowel movement

IMPORTANT SAFETY INFORMATION

MHRA/CHM ADVICE: CORTICOSTEROIDS: RARE RISK OF CENTRAL SEROUS CHORIOPATHY WITH LOCAL AS WELL AS SYSTEMIC ADMINISTRATION (AUGUST 2017)

See Corticosteroids, general use p. 450.

- **CAUTIONS** Local anaesthetic component can be absorbed through the rectal mucosa (avoid excessive application) · local anaesthetic component may cause sensitisation (use for short periods only—no longer than a few days)

- **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.

Ointment

- ▶ **Scheriproct** (Bayer Plc)
Prednisolone hexanoate 1.9 mg per 1 gram, Cinchocaine hydrochloride 5 mg per 1 gram Scheriproct ointment | 30 gram [PoM] £3.23 DT = £3.23

Suppository

- ▶ **Scheriproct** (Bayer Plc)
Cinchocaine hydrochloride 1 mg, Prednisolone hexanoate 1.3 mg Scheriproct suppositories | 12 suppository [PoM] £1.52 DT = £1.52

Hydrocortisone with lidocaine

21-Dec-2017

● INDICATIONS AND DOSE

Haemorrhoids | Pruritus ani

- ▶ BY RECTUM USING AEROSOL SPRAY
- ▶ Child 2–13 years (under medical advice only): 1 spray up to 3 times a day, spray once over the affected area
- ▶ Child 14–17 years: 1 spray up to 3 times a day for no longer than 7 days without medical advice, spray once over the affected area
- ▶ BY RECTUM USING OINTMENT
- ▶ Child: Apply several times daily, for short term use only

IMPORTANT SAFETY INFORMATION

MHRA/CHM ADVICE: CORTICOSTEROIDS: RARE RISK OF CENTRAL SEROUS CHORIOPATHY WITH LOCAL AS WELL AS SYSTEMIC ADMINISTRATION (AUGUST 2017)

See Corticosteroids, general use p. 450.

- **CAUTIONS** Local anaesthetic component can be absorbed through the rectal mucosa (avoid excessive application) · local anaesthetic component may cause sensitisation (use for short periods only—no longer than a few days)

- **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.

Ointment

- ▶ **Xyloproct** (Aspen Pharma Trading Ltd)
Hydrocortisone acetate 2.75 mg per 1 gram, Lidocaine 50 mg per 1 gram Xyloproct 5%/0.275% ointment | 20 gram [PoM] £4.19 DT = £4.19