

● SIDE-EFFECTS

- ▶ **Common or very common** Crying abnormal · drowsiness · eating disorder
- ▶ **Uncommon** Seizures · vascular disorders
- ▶ **Frequency not known** Extensive swelling of vaccinated limb · hypotonic-hyporesponsiveness episode

- **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.

Suspension for injection

EXCIPIENTS: May contain Kanamycin

- ▶ **Bexxero** (GlaxoSmithKline UK Ltd)
Bexxero vaccine suspension for injection 0.5ml pre-filled syringes | 1 pre-filled disposable injection [PoM] £75.00
- ▶ **Trumenba** (Pfizer Ltd) ▼
Trumenba vaccine suspension for injection 0.5ml pre-filled syringes | 1 pre-filled disposable injection [PoM] £75.00

F 814

Meningococcal group C vaccine

● INDICATIONS AND DOSE

Patients with confirmed serogroup C disease (who have previously been immunised)

- ▶ BY INTRAMUSCULAR INJECTION
- ▶ Child 1-17 years: 0.5 mL for 1 dose, dose to be given before discharge from hospital

● SIDE-EFFECTS

- ▶ **Common or very common** Anxiety · cough · crying · drowsiness · gastrointestinal discomfort · hyperhidrosis · increased risk of infection · pain · sleep disorders
- ▶ **Uncommon** Asthenia · chills · eyelid oedema · flushing · joint stiffness · musculoskeletal stiffness · nasal congestion · oedema · respiratory disorders · seizures · sensation abnormal · syncope
- ▶ **Rare or very rare** Circulatory collapse · influenza like illness
- ▶ **Frequency not known** Angioedema · apnoea · dyspnoea · hypotonic-hyporesponsiveness episode · immune thrombocytopenic purpura · meningism (but no evidence that vaccine causes meningococcal C meningitis) · Stevens-Johnson syndrome

- **PRESCRIBING AND DISPENSING INFORMATION** Available as part of childhood immunisation schedule from www.immform.dh.gov.uk.

- **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.

Suspension for injection

- ▶ **NeisVac-C** (Pfizer Ltd)
NeisVac-C vaccine suspension for injection 0.5ml pre-filled syringes | 10 pre-filled disposable injection [PoM] £187.50

F 814

Meningococcal groups A with C and W135 and Y vaccine

● INDICATIONS AND DOSE

MENVEO®

Primary immunisation against *Neisseria meningitidis*

- ▶ BY INTRAMUSCULAR INJECTION
- ▶ Child 13-15 years: 0.5 mL for 1 dose, dose preferably injected into deltoid region

Immunisation against *Neisseria meningitidis* in an unimmunised patient

- ▶ BY INTRAMUSCULAR INJECTION
- ▶ Child 10-17 years: 0.5 mL for 1 dose, booster dose is not required

Immunisation against *Neisseria meningitidis* in those at risk of exposure to prevent invasive disease

- ▶ BY INTRAMUSCULAR INJECTION
- ▶ Child 3-11 months: 0.5 mL every month for 2 doses, dose preferably injected into deltoid region
- ▶ Child 1-17 years: 0.5 mL for 1 dose, dose preferably injected into deltoid region

Patients attending university for the first time (who have not received the routine meningococcal groups A with C and W135 and Y conjugate vaccine over the age of 10 years)

- ▶ BY INTRAMUSCULAR INJECTION
- ▶ Child 16-17 years: 0.5 mL for 1 dose

NIMENRIX®

Primary immunisation against *Neisseria meningitidis*

- ▶ BY INTRAMUSCULAR INJECTION
- ▶ Child 13-15 years: 0.5 mL for 1 dose, to be injected preferably into deltoid region

Immunisation against *Neisseria meningitidis* in an unimmunised patient

- ▶ BY INTRAMUSCULAR INJECTION
- ▶ Child 10-17 years: 0.5 mL for 1 dose, booster dose is not required

Immunisation against *Neisseria meningitidis* in those at risk of exposure

- ▶ BY INTRAMUSCULAR INJECTION
- ▶ Child 1-17 years: 0.5 mL for 1 dose, to be injected preferably into deltoid region (or anterolateral thigh in child 12-23 months), then 0.5 mL after 1 year if required for 1 dose, second dose should be considered in those who continue to be at risk of *Neisseria meningitidis* serogroup A infection

Patients attending university for the first time (who have not received the routine meningococcal groups A with C and W135 and Y conjugate vaccine over the age of 10 years)

- ▶ BY INTRAMUSCULAR INJECTION
- ▶ Child 16-17 years: 0.5 mL for 1 dose

● UNLICENSED USE

MENVEO® *Menveo*® is not licensed for use in children under 2 years.

● SIDE-EFFECTS

- ▶ **Common or very common** Drowsiness
- ▶ **Uncommon** Crying · insomnia · numbness · pain in extremity
- ▶ **Frequency not known** Extensive swelling of vaccinated limb

- **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.

Powder and solvent for solution for injection

- ▶ **Menveo** (GlaxoSmithKline UK Ltd)
Menveo vaccine powder and solvent for solution for injection 0.5ml vials | 1 vial [PoM] £30.00
- ▶ **Nimenrix** (Pfizer Ltd)
Nimenrix vaccine powder and solvent for solution for injection 0.5ml pre-filled syringes | 1 pre-filled disposable injection [PoM] £30.00 DT = £30.00

F 814

Pneumococcal polysaccharide conjugate vaccine (adsorbed)

01-May-2019

● INDICATIONS AND DOSE

PREVENAR 13®

Primary immunisation against pneumococcal infection [first dose]

- ▶ BY INTRAMUSCULAR INJECTION
- ▶ Child 2 months: 0.5 mL for 1 dose, anterolateral thigh is preferred site of injection in infants under 1 year

continued →