

guideline: **Acute Kidney Injury** (www.nice.org.uk/guidance/ng148).

Glomerular filtration rate is low at birth and increases rapidly during the first 6 months. Thereafter, glomerular filtration rate increases gradually to reach adult levels by 1–2 years of age, when standardised to a typical adult body surface area (1.73 m²). In the first weeks after birth, serum creatinine falls; a single measure of serum creatinine provides only a crude estimate of renal function and observing the change over days is of more use. In the neonate, a sustained rise in serum creatinine or a lack of the expected postnatal decline, is indicative of a reduced glomerular filtration rate.

Dose recommendations are based on the severity of renal impairment. This is expressed in terms of **glomerular filtration rate** (mL/minute/1.73 m²).

The following equations provide a guide to glomerular filtration rate.

Child over 1 year:

$$\text{Estimated glomerular filtration rate (mL/minute/1.73 m}^2\text{)} = 40 \times \text{height (cm)/serum creatinine (micromol/litre)}$$

Neonate:

$$\text{Estimated glomerular filtration rate (mL/minute/1.73 m}^2\text{)} = 30 \times \text{height (cm)/serum creatinine (micromol/litre)}$$

The values used in these formulas may differ according to locality or laboratory.

The serum-creatinine concentration is sometimes used as a measure of renal function but is only a **rough guide** even when corrected for age, weight, and sex.

Important The information on dose adjustment in *BNF for Children* is expressed in terms of estimated glomerular filtration rate. Renal function in adults is increasingly being reported as estimated glomerular filtration rate (eGFR) normalised to a body surface area of 1.73 m²; however, eGFR is derived from the MDRD (Modification of Diet in Renal Disease) formula which is not validated for use in children. eGFR derived from the MDRD formula should **not** be used to adjust drug doses in children with renal impairment.

In *BNF for Children*, values for measures of renal function are included where possible. However, where such values are not available, the *BNF for Children* reflects the terms used in the published information.

Degrees of renal impairment defined using estimated glomerular filtration rate (eGFR)

Chronic kidney disease in adults: UK guidelines for identification, management and referral (March 2006) defines renal function as follows:

Degree of impairment	eGFR ¹ mL/minute/1.73 m ²
Normal: Stage 1	More than 90 (with other evidence of kidney damage)
Mild: Stage 2	60–89 (with other evidence of kidney damage)
Moderate ² : Stage 3	30–59
Severe: Stage 4	15–29
Established renal failure: Stage 5	Less than 15

1. Estimated glomerular filtration rate (eGFR) derived from the Modification of Diet in Renal Disease (MDRD) formula for use in patients over 18 years
2. NICE clinical guideline 73 (September 2008)–Chronic kidney disease: Stage 3A eGFR = 45–59, Stage 3B eGFR = 30–44

If even mild renal impairment is considered likely on clinical grounds, renal function should be checked before prescribing **any** drug which requires dose modification.

Where care is needed when prescribing in renal impairment, this is indicated under the relevant drug in *BNF for Children*.

Dialysis

For prescribing in children on renal replacement therapy consult specialist literature.

Advanced Pharmacy Services

Children with renal impairment may be eligible for the Medicines Use Review service provided by a community pharmacist. For further information, see *Advanced Pharmacy Services* in Medicines optimisation p. 25.

Drug prescribing should be kept to the minimum in all children with severe renal disease.