

- **CAUTIONS** Local anaesthetic component can be absorbed through the rectal mucosa (avoid excessive application, particularly in children and infants) · local anaesthetic component may cause sensitisation (use for short periods only—no longer than a few days)
- **PRESCRIBING AND DISPENSING INFORMATION** A proprietary brand *Anusol Plus HC*® (ointment and suppositories) is on sale to the public.
- **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.

Ointment

- ▶ **Anusol-HC** (Church & Dwight UK Ltd)
Hydrocortisone acetate 2.5 mg per 1 gram, Bismuth oxide 8.75 mg per 1 gram, Benzyl benzoate 12.5 mg per 1 gram, Peru Balsam 18.75 mg per 1 gram, Bismuth subgallate 22.5 mg per 1 gram, Zinc oxide 107.5 mg per 1 gram Anusol HC ointment | 30 gram [PoM] £2.49

Suppository

- ▶ **Anusol-HC** (Church & Dwight UK Ltd)
Hydrocortisone acetate 10 mg, Bismuth oxide 24 mg, Benzyl benzoate 33 mg, Peru Balsam 49 mg, Bismuth subgallate 59 mg, Zinc oxide 296 mg Anusol HC suppositories | 12 suppository [PoM] £1.74

Cinchocaine hydrochloride with fluocortolone caproate and fluocortolone pivalate

21-Dec-2017

INDICATIONS AND DOSE**Haemorrhoids | Pruritus ani**

- ▶ BY RECTUM USING OINTMENT
- ▶ Child: Apply twice daily for 5–7 days, apply 3–4 times a day if required, on the first day of treatment, then apply once daily for a few days after symptoms have cleared
- ▶ BY RECTUM USING SUPPOSITORIES
- ▶ Child 12–17 years: Initially 1 suppository daily for 5–7 days, to be inserted after a bowel movement, then 1 suppository once daily on alternate days for 1 week

Haemorrhoids (severe cases) | Pruritus ani (severe cases)

- ▶ BY RECTUM USING SUPPOSITORIES
- ▶ Child 12–17 years: Initially 1 suppository 2–3 times a day for 5–7 days, then 1 suppository once daily on alternate days for 1 week

IMPORTANT SAFETY INFORMATION

MHRA/CHM ADVICE: CORTICOSTEROIDS: RARE RISK OF CENTRAL SEROUS CHORIORETINOPATHY WITH LOCAL AS WELL AS SYSTEMIC ADMINISTRATION (AUGUST 2017)
See Corticosteroids, general use p. 470.

- **CAUTIONS** Local anaesthetic component can be absorbed through the rectal mucosa (avoid excessive application) · local anaesthetic component may cause sensitisation (use for short periods only—no longer than a few days)
- **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.

Ointment

- ▶ **Ultraproct** (Meadow Laboratories Ltd)
Fluocortolone pivalate 920 microgram per 1 gram, Fluocortolone caproate 950 microgram per 1 gram, Cinchocaine hydrochloride 5 mg per 1 gram Ultraproct ointment | 30 gram [PoM] £8.27

Suppository

- ▶ **Ultraproct** (Meadow Laboratories Ltd)
Fluocortolone pivalate 610 microgram, Fluocortolone caproate 630 microgram, Cinchocaine hydrochloride 1 mg Ultraproct suppositories | 12 suppository [PoM] £4.06

Cinchocaine with hydrocortisone

21-Dec-2017

INDICATIONS AND DOSE**PROCTOSEDYL® OINTMENT****Haemorrhoids | Pruritus ani**

- ▶ TO THE SKIN, OR BY RECTUM
- ▶ Child: Apply twice daily, to be administered morning and night and after a bowel movement. Apply externally or by rectum. Do not use for longer than 7 days

PROCTOSEDYL® SUPPOSITORIES**Haemorrhoids | Pruritus ani**

- ▶ BY RECTUM
- ▶ Child 12–17 years: 1 suppository, insert suppository night and morning and after a bowel movement. Do not use for longer than 7 days

UNIROID-HC® OINTMENT**Haemorrhoids | Pruritus ani**

- ▶ TO THE SKIN, OR BY RECTUM
- ▶ Child 1 month–11 years (under medical advice only): Apply twice daily, and apply after a bowel movement, apply externally or by rectum, do not use for longer than 7 days
- ▶ Child 12–17 years: Apply twice daily, and apply after a bowel movement, apply externally or by rectum, do not use for longer than 7 days

UNIROID-HC® SUPPOSITORIES**Haemorrhoids | Pruritus ani**

- ▶ BY RECTUM
- ▶ Child 12–17 years: 1 suppository, insert twice daily and after a bowel movement. Do not use for longer than 7 days

IMPORTANT SAFETY INFORMATION

MHRA/CHM ADVICE: CORTICOSTEROIDS: RARE RISK OF CENTRAL SEROUS CHORIORETINOPATHY WITH LOCAL AS WELL AS SYSTEMIC ADMINISTRATION (AUGUST 2017)
See Corticosteroids, general use p. 470.

- **CAUTIONS** Local anaesthetic component can be absorbed through the rectal mucosa (avoid excessive application, particularly in children and infants) · local anaesthetic component may cause sensitisation (use for short periods only—no longer than a few days)
 - **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.
- Ointment**
- ▶ **Proctosedyl** (Sanofi)
Cinchocaine hydrochloride 5 mg per 1 gram, Hydrocortisone 5 mg per 1 gram Proctosedyl ointment | 30 gram [PoM] £10.34 DT = £10.34
 - ▶ **Uniroid HC** (Chemidex Pharma Ltd)
Cinchocaine hydrochloride 5 mg per 1 gram, Hydrocortisone 5 mg per 1 gram Uniroid HC ointment | 30 gram [PoM] £4.23 DT = £10.34
- Suppository**
- ▶ **Proctosedyl** (Sanofi)
Cinchocaine hydrochloride 5 mg, Hydrocortisone 5 mg Proctosedyl suppositories | 12 suppository [PoM] £5.08 DT = £5.08
 - ▶ **Uniroid HC** (Chemidex Pharma Ltd)
Cinchocaine hydrochloride 5 mg, Hydrocortisone 5 mg Uniroid HC suppositories | 12 suppository [PoM] £1.91 DT = £5.08