

hypertriglyceridaemia · inflammatory bowel disease · menstrual cycle irregularities · suicidal tendencies

- **PREGNANCY** Avoid—risk of feminisation of male fetus with cyproterone.
- **BREAST FEEDING** Manufacturer advises avoid; possibility of anti-androgen effects in neonate with cyproterone.
- **HEPATIC IMPAIRMENT** Manufacturer advises caution; avoid in severe or active disease
- **PRESCRIBING AND DISPENSING INFORMATION** A mixture of cyproterone acetate and ethinylestradiol in the mass proportions 2000 parts to 35 parts, respectively.

- **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.

Tablet

▶ Co-cyprindiol (Non-proprietary)

Ethinylestradiol 35 microgram, Cyproterone acetate 2 mg Co-cyprindiol 2000microgram/35microgram tablets | 63 tablet [PoM] £16.65 DT = £10.78

▶ Clairette (Stragen UK Ltd)

Ethinylestradiol 35 microgram, Cyproterone acetate 2 mg Clairette 2000/35 tablets | 63 tablet [PoM] £5.90 DT = £10.78

▶ Dianette (Bayer Plc)

Ethinylestradiol 35 microgram, Cyproterone acetate 2 mg Dianette tablets | 63 tablet [PoM] £7.71 DT = £10.78

▶ Teragezza (Morningside Healthcare Ltd)

Ethinylestradiol 35 microgram, Cyproterone acetate 2 mg Teragezza 2000microgram/35microgram tablets | 63 tablet [PoM] £11.10 DT = £10.78

ANTIBACTERIALS > LINCOSAMIDES

Clindamycin

18-May-2020

● INDICATIONS AND DOSE

DALACIN T[®] LOTION

Acne vulgaris

- ▶ TO THE SKIN
- ▶ Child: Apply twice daily, to be applied thinly

DALACIN T[®] SOLUTION

Acne vulgaris

- ▶ TO THE SKIN
- ▶ Child: Apply twice daily, to be applied thinly

ZINDACLIN[®] GEL

Acne vulgaris

- ▶ TO THE SKIN
- ▶ Child 12-17 years: Apply once daily, to be applied thinly

- **INTERACTIONS** → Appendix 1: clindamycin
- **SIDE-EFFECTS**
 - ▶ Common or very common Skin reactions
 - ▶ Frequency not known Abdominal pain · antibiotic associated colitis · folliculitis gram-negative · gastrointestinal disorder
- **PATIENT AND CARER ADVICE** Patients and their carers should be advised to discontinue and contact a doctor immediately if severe, prolonged or bloody diarrhoea develops.
- **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.

Gel

EXCIPIENTS: May contain Propylene glycol

▶ Zindaclin (Crawford Healthcare Ltd)

Clindamycin (as Clindamycin phosphate) 10 mg per 1 gram Zindaclin 1% gel | 30 gram [PoM] £8.66 DT = £8.66

Liquid

EXCIPIENTS: May contain Cetostearyl alcohol (including cetyl and stearyl alcohol), hydroxybenzoates (parabens), propylene glycol

▶ Dalacin T (Pfizer Ltd)

Clindamycin (as Clindamycin phosphate) 10 mg per 1 ml Dalacin T 1% topical lotion | 30 ml [PoM] £5.08 DT = £5.08 | 60 ml [PoM] £10.16
Dalacin T 1% topical solution | 30 ml [PoM] £4.34 DT = £4.34 | 50 ml [PoM] £7.23

Combinations available: *Benzoyl peroxide with clindamycin*, p. 808 · *Tretinoin with clindamycin*, p. 810

ANTIBACTERIALS > MACROLIDES

Erythromycin with zinc acetate

20-Apr-2020

The properties listed below are those particular to the combination only. For the properties of the components please consider, erythromycin p. 355.

● INDICATIONS AND DOSE

Acne vulgaris

- ▶ TO THE SKIN
- ▶ Child: Apply twice daily

- **INTERACTIONS** → Appendix 1: macrolides

- **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.

Liquid

▶ Zineryt (LEO Pharma)

Zinc acetate 12 mg per 1 ml, Erythromycin 40 mg per 1 ml Zineryt lotion | 30 ml [PoM] £9.25 DT = £9.25 | 90 ml [PoM] £20.02 DT = £20.02

ANTISEPTICS AND DISINFECTANTS > PEROXIDES

Benzoyl peroxide

● INDICATIONS AND DOSE

Acne vulgaris

- ▶ TO THE SKIN
- ▶ Child 12-17 years: Apply 1–2 times a day, preferably apply after washing with soap and water, start treatment with lower-strength preparations

Infantile acne

- ▶ TO THE SKIN
- ▶ Child 1 month-1 year: Apply 1–2 times a day, start treatment with lower-strength preparations

- **UNLICENSED USE** Not licensed for use in treatment of infantile acne.
- **CAUTIONS** Avoid contact with broken skin · avoid contact with eyes · avoid contact with mouth · avoid contact with mucous membranes · avoid excessive exposure to sunlight
- **SIDE-EFFECTS**
 - ▶ Common or very common Skin reactions
 - ▶ Frequency not known Facial swelling
- **SIDE-EFFECTS, FURTHER INFORMATION** Reduce frequency or suspend use until skin irritation subsides and re-introduce at reduced frequency.
- **PATIENT AND CARER ADVICE** May bleach fabrics and hair.
- **MEDICINAL FORMS** There can be variation in the licensing of different medicines containing the same drug.

Gel

EXCIPIENTS: May contain Fragrances, propylene glycol

▶ Acnecide (Galderma (UK) Ltd)

Benzoyl peroxide 50 mg per 1 gram Acnecide 5% gel | 30 gram [P] £5.44 DT = £5.44 | 60 gram [P] £10.68 DT = £10.68
Acnecide Wash 5% gel | 50 gram [P] £5.44 DT = £5.44

Combinations available: *Adapalene with benzoyl peroxide*, p. 808