

1 Anaesthesia adjuvants

Pre-medication and peri-operative drugs

Drugs that affect gastric pH

Regurgitation and aspiration of gastric contents (Mendelson's syndrome) can be a complication of general anaesthesia, particularly in obstetrics and in gastro-oesophageal reflux disease; prophylaxis against acid aspiration is not routinely used in children but may be required in high-risk cases.

An **H₂-receptor antagonist** can be used before surgery to increase the pH and reduce the volume of gastric fluid. It does not affect the pH of fluid already in the stomach and this limits its value in emergency procedures; an oral H₂-receptor antagonist can be given 1–2 hours before the procedure.

Antimuscarinic drugs

Antimuscarinic drugs are used (less commonly nowadays) as premedicants to dry bronchial and salivary secretions which are increased by intubation, upper airway surgery, or some inhalational anaesthetics. They are also used before or with neostigmine p. 700 to prevent bradycardia, excessive salivation, and other muscarinic actions of neostigmine. They also prevent bradycardia and hypotension associated with drugs such as propofol p. 864 and suxamethonium chloride p. 872.

Atropine sulfate p. 869 is now rarely used for premedication but still has an emergency role in the treatment of vagotonic side-effects. Atropine sulfate may have a role in cardiopulmonary resuscitation.

Hyoscine hydrobromide p. 283 reduces secretions and also provides a degree of amnesia, sedation, and anti-emesis. Unlike atropine sulfate it may produce bradycardia rather than tachycardia.

Glycopyrronium bromide p. 870 reduces salivary secretions. When given intravenously it produces less tachycardia than atropine sulfate. It is widely used with neostigmine for reversal of non-depolarising muscle relaxants.

Glycopyrronium bromide or hyoscine hydrobromide are also used to control excessive secretions in upper airways or hypersalivation in palliative care and in children unable to control posture or with abnormal swallowing reflex; effective dose varies and tolerance may develop. The intramuscular route should be avoided if possible. Hyoscine hydrobromide transdermal patches may also be used.

Sedative drugs

Premedication

Fear and anxiety before a procedure (including the night before) can be minimised by using a sedative drug, usually a **benzodiazepine**. Premedication may also augment the action of anaesthetics and provide some degree of pre-operative amnesia. The choice of drug depends on the individual, the nature of the procedure, the anaesthetic to be used, and other prevailing circumstances such as outpatients, obstetrics, and availability of recovery facilities. The choice also varies between elective and emergency procedures. Oral administration is preferred if possible; the rectal route should only be used in exceptional circumstances.

Premedicants can be given the night before major surgery; a further, smaller dose may be required before surgery. Alternatively, the first dose may be given on the day of the procedure.

Oral midazolam p. 239 is the most common premedicant for children; temazepam p. 879 may be used in older children. The antihistamine alimemazine tartrate p. 186 is occasionally used orally, but when given alone it may cause postoperative restlessness in the presence of pain.

Benzodiazepines

Benzodiazepines possess useful properties for premedication including relief of anxiety, sedation, and amnesia; short-acting benzodiazepines taken by mouth are the most common premedicants. Benzodiazepines are also used for sedation prior to clinical procedures and for sedation in intensive care.

Benzodiazepines may occasionally cause marked respiratory depression and facilities for its treatment are essential; flumazenil p. 899 is used to antagonise the effects of benzodiazepines.

Midazolam, a water-soluble benzodiazepine, is the preferred benzodiazepine for premedication and for sedation for clinical procedures in children. It has a fast onset of action, and recovery is faster than for other benzodiazepines. Recovery may be longer in children with a low cardiac output, or after repeated dosing.

Midazolam can be given by mouth [unlicensed], but its bitter acidic taste may need to be disguised. It can also be given buccally [unlicensed indication] or intranasally [unlicensed]. Midazolam is associated with profound sedation when high doses are given or when it is used with certain other drugs. It can cause severe disinhibition and restlessness in some children. Midazolam is not recommended for prolonged sedation in neonates; drug accumulation is likely to occur.

Temazepam is given by mouth for premedication in older children and has a short duration of action. Anxiolytic and sedative effects last about 90 minutes, although there may be residual drowsiness. Temazepam is rarely used for dental procedures in children.

Lorazepam p. 238 produces more prolonged sedation than temazepam and it has marked amnesic effects.

Peri-operative use of diazepam p. 236 is not recommended in children; onset and magnitude of response are unreliable, and paradoxical effects may occur. Diazepam is not used for dental procedures in children.

Antagonists for central and respiratory depression

Respiratory depression is a major concern with opioid analgesics and it may be treated by artificial ventilation or be reversed by an opioid antagonist. Naloxone hydrochloride p. 901 given intravenously immediately reverses opioid-induced respiratory depression but the dose may have to be repeated because of its **short duration of action**.

Intramuscular injection of naloxone hydrochloride produces a more gradual and prolonged effect but absorption may be erratic. Care is required in children requiring pain relief because naloxone hydrochloride also antagonises the analgesic effect of opioids.

Flumazenil is a benzodiazepine antagonist for the reversal of the central sedative effects of benzodiazepines after anaesthetic and similar procedures. Flumazenil has a shorter half-life and duration of action than diazepam or midazolam so patients may become resedated.

Neonates

Naloxone hydrochloride is used in newborn infants to reverse respiratory depression and sedation resulting from the use of opioids by the mother, usually for pain during labour. In neonates the effects of opioids may persist for up to 48 hours and in such cases naloxone hydrochloride is often given by intramuscular injection for its prolonged effect. In severe respiratory depression after birth, breathing should first be established (using artificial means if necessary) and naloxone hydrochloride administered only if use of opioids by the mother is thought to cause the respiratory depression; the infant should be monitored