

following cardiac surgery. It is given by continuous 24-hour intravenous infusion.

Epoprostenol is a powerful inhibitor of platelet aggregation and there is a possible risk of haemorrhage. It is sometimes used as an antiplatelet in renal dialysis when heparins are unsuitable or contra-indicated. It can also cause serious systemic hypotension and, if withdrawn suddenly, can cause pulmonary hypertensive crisis.

Children on prolonged treatment can become tolerant to epoprostenol, and therefore require an increase in dose.

Iloprost p. 127 is a synthetic analogue of epoprostenol and is efficacious when nebulised in adults with pulmonary arterial hypertension, but experience in children is limited. It is more stable than epoprostenol and has a longer half-life.

Bosentan p. 125 is a dual endothelin receptor antagonist used orally in the treatment of pulmonary arterial hypertension. The concentration of endothelin, a potent vasoconstrictor, is raised in sustained pulmonary hypertension.

Sildenafil, a vasodilator developed for the treatment of erectile dysfunction, is also used for pulmonary arterial hypertension. It is used either alone or as an adjunct to other drugs.

Sildenafil is a selective phosphodiesterase type-5 inhibitor. Inhibition of this enzyme in the lungs enhances the vasodilatory effects of nitric oxide and promotes relaxation of vascular smooth muscle.

Sildenafil has also been used in pulmonary hypertension for weaning children off inhaled nitric oxide following cardiac surgery, and less successfully in idiopathic pulmonary arterial hypertension.

Tolazoline p. 128 is now rarely used to correct pulmonary artery vasospasm in pulmonary hypertension of the newborn as better alternatives are available. Tolazoline is an alpha-blocker and produces both pulmonary and systemic vasodilation.

Advanced Pharmacy Services

Children with hypertension may be eligible for the New Medicines Service / Medicines Use Review service provided by a community pharmacist. For further information, see *Advanced Pharmacy Services* in Medicines optimisation p. 25.

Antihypertensive drugs

Vasodilator antihypertensive drugs

Vasodilators have a potent hypotensive effect, especially when used in combination with a beta-blocker and a thiazide. **Important:** see Hypertension (hypertensive emergencies) for a warning on the hazards of a very rapid fall in blood pressure.

Hydralazine hydrochloride p. 123 is given by mouth as an adjunct to other antihypertensives for the treatment of resistant hypertension but is rarely used; when used alone it causes tachycardia and fluid retention.

Sodium nitroprusside p. 124 is given by intravenous infusion to control severe hypertensive crisis when parenteral treatment is necessary. At low doses it reduces systemic vascular resistance and increases cardiac output; at high doses it can produce profound systemic hypotension—continuous blood pressure monitoring is therefore essential. Sodium nitroprusside may also be used to control paradoxical hypertension after surgery for coarctation of the aorta.

Minoxidil p. 124 should be reserved for the treatment of severe hypertension resistant to other drugs. Vasodilatation is accompanied by increased cardiac output and tachycardia and children develop fluid retention. For this reason the addition of a beta-blocker and a diuretic (usually furosemide p. 148, in high dosage) are mandatory. Hypertrichosis is troublesome and renders this drug unsuitable for females.

Prazosin p. 107 and doxazosin p. 529 have alpha-blocking and vasodilator properties.

Centrally acting antihypertensive drugs

Methyldopa, a centrally acting antihypertensive, is of little value in the management of refractory sustained hypertension in infants and children. On prolonged use it is associated with fluid retention (which may be alleviated by concomitant use of diuretics).

Methyldopa is also effective for the management of hypertension in pregnancy.

Clonidine hydrochloride p. 108 is also a centrally acting antihypertensive but has the disadvantage that sudden withdrawal may cause a hypertensive crisis. Clonidine hydrochloride is also used under specialist supervision for pain management, sedation, and opioid withdrawal, attention deficit hyperactivity disorder, and Tourette syndrome.

Adrenergic neurone blocking drugs

Adrenergic neurone blocking drugs prevent the release of noradrenaline from postganglionic adrenergic neurones. These drugs do not control supine blood pressure and may cause postural hypotension. For this reason they have largely fallen from use in adults and are rarely used in children.

Alpha-adrenoceptor blocking drugs

Doxazosin and prazosin have post-synaptic alpha-blocking and vasodilator properties and rarely cause tachycardia. They can, however, reduce blood pressure rapidly after the first dose and should be introduced with caution.

Alpha-blockers can be used with other antihypertensive drugs in the treatment of resistant hypertension.

Drugs affecting the renin-angiotensin system

Angiotensin-converting enzyme inhibitors

Angiotensin-converting enzyme inhibitors (ACE inhibitors) inhibit the conversion of angiotensin I to angiotensin II. The main indications of ACE inhibitors in children are shown below. In infants and young children, captopril p. 118 is often considered first.

Initiation under specialist supervision

Treatment with ACE inhibitors should be initiated under specialist supervision and with careful clinical monitoring in children.

Heart failure

ACE inhibitors have a valuable role in all grades of heart failure, usually combined with a loop diuretic. Potassium supplements and potassium-sparing diuretics should be discontinued before introducing an ACE inhibitor because of the risk of hyperkalaemia. Profound first-dose hypotension can occur when ACE inhibitors are introduced to children with heart failure who are already taking a high dose of a loop diuretic. Temporary withdrawal of the loop diuretic reduces the risk, but can cause severe rebound pulmonary oedema.

Hypertension

ACE inhibitors may be considered for hypertension when thiazides and beta-blockers are contra-indicated, not tolerated, or fail to control blood pressure; they may be considered for hypertension in children with type 1 diabetes with nephropathy. ACE inhibitors can reduce blood pressure very rapidly in some patients particularly in those receiving diuretic therapy.