

Chapter 14

Vaccines

CONTENTS

1 Immunoglobulin therapy	page 821	3 Tuberculosis diagnostic test	page 828
2 Post-exposure prophylaxis	827	4 Vaccination	828

1 Immunoglobulin therapy

IMMUNE SERA AND IMMUNOGLOBULINS > IMMUNOGLOBULINS

Immunoglobulins

04-Oct-2017

Passive immunity

Immunity with immediate protection against certain infective organisms can be obtained by injecting preparations made from the plasma of immune individuals with adequate levels of antibody to the disease for which protection is sought. The duration of this passive immunity varies according to the dose and the type of immunoglobulin. Passive immunity may last only a few weeks; when necessary, passive immunisation can be repeated. Antibodies of human origin are usually termed immunoglobulins. The term antiserum is applied to material prepared in animals. Because of serum sickness and other allergic-type reactions that may follow injections of antiserum, this therapy has been replaced wherever possible by the use of immunoglobulins. Reactions are theoretically possible after injection of human immunoglobulins but reports of such reactions are very rare.

Two types of human immunoglobulin preparation are available, normal immunoglobulin p. 825 and **disease-specific immunoglobulins**.

Human immunoglobulin is a sterile preparation of concentrated antibodies (immune globulins) recovered from pooled human plasma or serum obtained from outside the UK, tested and found non-reactive for hepatitis B surface antigen and for antibodies against hepatitis C virus and human immunodeficiency virus (types 1 and 2). A global shortage of human immunoglobulin and the rapidly increasing range of clinical indications for treatment with immunoglobulins has resulted in the need for a Demand Management programme in the UK, for further information consult www.ivig.nhs.uk and *Clinical Guidelines for Immunoglobulin Use*, www.gov.uk/dh.

Further information on the use of immunoglobulins is included in Public Health England's *Immunoglobulin Handbook*, www.gov.uk/phe, and in the Department of Health's publication, *Immunisation against Infectious Disease*, www.gov.uk/dh.

Availability

Normal immunoglobulin for intramuscular administration is available from some regional Public Health laboratories for protection of contacts and the control of outbreaks of hepatitis A, measles, and rubella only. For other indications, subcutaneous or intravenous normal immunoglobulin should be purchased from the manufacturer.

Disease-specific immunoglobulins are available from some regional Public Health laboratories, with the exception of tetanus immunoglobulin p. 826 which is available from BPL,

hospital pharmacies, or blood transfusion departments. Rabies immunoglobulin p. 826 is available from the Specialist and Reference Microbiology Division, Public Health England, Colindale. Hepatitis B immunoglobulin p. 824 required by transplant centres should be obtained commercially.

In Scotland all immunoglobulins are available from the *Scottish National Blood Transfusion Service (SNBTS)*.

In Wales all immunoglobulins are available from the *Welsh Blood Service (WBS)*.

In Northern Ireland all immunoglobulins are available from the *Northern Ireland Blood Transfusion Service (NI BTS)*.

Normal immunoglobulin

Human normal immunoglobulin ('HNIG') is prepared from pools of at least 1000 donations of human plasma; it contains immunoglobulin G (IgG) and antibodies to hepatitis A, measles, mumps, rubella, varicella, and other viruses that are currently prevalent in the general population.

Uses

Normal immunoglobulin (containing 10–18% protein) is administered by **intramuscular injection** for the protection of susceptible contacts against **hepatitis A virus** (infectious hepatitis), **measles** and, to a lesser extent, **rubella**. Injection of immunoglobulin produces immediate protection lasting for several weeks.

Normal immunoglobulin (containing 3–12% protein) for **intravenous administration** is used as **replacement therapy** for children with congenital agammaglobulinaemia and hypogammaglobulinaemia, and for the short-term treatment of idiopathic thrombocytopenic purpura and Kawasaki disease; it is also used for the prophylaxis of infection following bone-marrow transplantation and in children with symptomatic HIV infection who have recurrent bacterial infections. Normal immunoglobulin for replacement therapy may also be given intramuscularly or subcutaneously, but intravenous formulations are normally preferred. Intravenous immunoglobulin is also used in the treatment of Guillain-Barré syndrome as an alternative to plasma exchange.

The dose of normal immunoglobulin used as replacement therapy in patients with immunodeficiencies is **not the same** as the dose required for treatment of acute conditions. For Kawasaki disease a single dose by intravenous infusion should be given with concomitant aspirin p. 98 within 10 days of onset of symptoms (but children with a delayed diagnosis may also benefit).

For guidance on the use of intravenous normal immunoglobulin and alternative therapies for other conditions, consult *Clinical Guidelines for Immunoglobulin Use* (www.gov.uk/dh).

Hepatitis A

Hepatitis A vaccine p. 852 is recommended for individuals at risk of infection including those visiting areas where the disease is highly endemic (all countries excluding Northern and Western Europe, North America, Japan, Australia, and