

Chapter 16

Emergency treatment of poisoning

CONTENTS

1 Active elimination from the gastrointestinal tract	page 898	3.2 Digoxin toxicity	page 900
2 Chemical toxicity	898	3.3 Heparin toxicity	900
2.1 Cyanide toxicity	898	3.4 Opioid toxicity	901
2.2 Organophosphorus toxicity	899	3.5 Paracetamol toxicity	902
3 Drug toxicity	899	4 Methaemoglobinemia	903
3.1 Benzodiazepine toxicity	899	5 Snake bites	904

Poisoning, emergency treatment

28-Apr-2020

Overview

These notes provide only an overview of the treatment of poisoning, and it is strongly recommended that either **TOXBASE** or the **UK National Poisons Information Service** be consulted when there is doubt about the degree of risk or about management.

Most childhood poisoning is accidental. Other causes include intentional overdose, drug abuse, iatrogenic and deliberate poisoning. The drugs most commonly involved in childhood poisoning are paracetamol p. 288, ibuprofen p. 707, orally ingested creams, aspirin p. 98, iron preparations, cough medicines, and the contraceptive pill.

Hospital admission

Children who have features of poisoning should generally be admitted to hospital. Children who have taken poisons with delayed actions should also be admitted, even if they appear well. Delayed-action poisons include aspirin, iron, paracetamol, tricyclic antidepressants, and co-phenotrope (diphenoxylate with atropine, *Lomotil*[®]) p. 53; the effects of modified-release preparations are also delayed. A note of all relevant information, including what treatment has been given, should accompany the patient to hospital.

Further information

TOXBASE, the primary clinical toxicology database of the National Poisons Information Service, is available on the internet to registered users at www.toxbase.org (a backup site is available at www.toxbasebackup.org if the main site cannot be accessed). It provides information about routine diagnosis, treatment, and management of patients exposed to drugs, household products, and industrial and agricultural chemicals.

Specialist information and advice on the treatment of poisoning is available day and night from the **UK National Poisons Information Service** on the following number: Tel: 0344 892 0111.

Advice on laboratory analytical services can be obtained from TOXBASE or from the National Poisons Information Service. Help with identifying capsules or tablets may be available from a regional medicines information centre or from the National Poisons Information Service (out of hours).

General care

It is often impossible to establish with certainty the identity of the poison and the size of the dose. This is not usually important because only a few poisons (such as opioids,

paracetamol, and iron) have specific antidotes; few patients require active removal of the poison. In most patients, treatment is directed at managing symptoms as they arise. Nevertheless, knowledge of the type and timing of poisoning can help in anticipating the course of events. All relevant information should be sought from the poisoned individual and from carers or parents. However, such information should be interpreted with care because it may not be complete or entirely reliable. Sometimes symptoms arise from other illnesses and patients should be assessed carefully. Accidents may involve domestic and industrial products (the contents of which are not generally known). The **National Poisons Information Service** should be consulted when there is doubt about any aspect of suspected poisoning.

Respiration

Respiration is often impaired in unconscious patients. An obstructed airway requires immediate attention. In the absence of trauma, the airway should be opened with simple measures such as chin lift or jaw thrust. An oropharyngeal or nasopharyngeal airway may be useful in patients with reduced consciousness to prevent obstruction, provided ventilation is adequate. Intubation and ventilation should be considered in patients whose airway cannot be protected or who have respiratory acidosis because of inadequate ventilation; such patients should be monitored in a critical care area.

Most poisons that impair consciousness also depress respiration. Assisted ventilation (either mouth-to-mouth or using a bag-valve-mask device) may be needed. Oxygen is not a substitute for adequate ventilation, although it should be given in the highest concentration possible in poisoning with carbon monoxide and irritant gases.

The potential for pulmonary aspiration of gastric contents should be considered.

Blood pressure

Hypotension is common in severe poisoning with central nervous system depressants; if severe, this may lead to irreversible brain damage or renal tubular necrosis. Hypotension should be corrected initially by raising the foot of the bed and administration of an infusion of either sodium chloride p. 636 or a colloid. Vasoconstrictor sympathomimetics are rarely required and their use may be discussed with the National Poisons Information Service or a paediatric intensive care unit.

Fluid depletion without hypotension is common after prolonged coma and after aspirin poisoning due to vomiting, sweating, and hyperpnoea.