

Chapter 11

Eye

CONTENTS

1 Allergic and inflammatory eye conditions	page 718
1.1 Allergic conjunctivitis	718
1.2 Inflammatory eye conditions	720
1.2a Anterior uveitis	722
2 Dry eye conditions	723
3 Eye infections	726
3.1 Bacterial eye infection	727
3.2 Viral eye infection	730

3.2a Ophthalmic herpes simplex	page 730
4 Eye procedures	730
4.1 Post-operative pain and inflammation	731
5 Glaucoma and ocular hypertension	733
6 Retinal disorders	739
6.1 Optic neuropathy	739

Eye

Eye treatment: drug administration

Drugs are most commonly administered to the eye by topical application as eye drops or eye ointments. When a higher drug concentration is required within the eye, a local injection may be necessary.

Eye-drop dispenser devices are available to aid the instillation of eye drops from plastic bottles and some are prescribable on the NHS (consult Drug Tariff—see Appliances and Reagents). Product-specific devices may be supplied by manufacturers—contact individual manufacturers for further information. They are particularly useful for children in whom normal application is difficult, for the visually impaired, or otherwise physically limited patients.

Eye drops and eye ointments

Eye drops are generally instilled into the pocket formed by gently pulling down the lower eyelid and keeping the eye closed for as long as possible after application; in neonates and infants it may be more appropriate to administer the drop in the inner angle of the open eye. One drop is all that is needed; instillation of more than one drop at a time should be discouraged because it may increase systemic side-effects. A small amount of eye ointment is applied similarly; the ointment melts rapidly and blinking helps to spread it.

When two different eye-drop preparations are used at the same time of day, dilution and overflow can occur when one immediately follows the other. The carer or child should therefore leave an interval of at least 5 minutes between the two; the interval should be extended when eye drops with a prolonged contact time, such as gels and suspensions, are used. Eye ointment should be applied after drops. Both drops and ointment can cause transient blurred vision; children should be warned, where appropriate, not to perform skilled tasks (e.g. cycling or driving) until vision is clear.

Systemic effects may arise from absorption of drugs into the general circulation from conjunctival vessels or from the nasal mucosa after the excess preparation has drained down through the tear ducts. The extent of systemic absorption following ocular administration is highly variable; nasal drainage of drugs is associated with eye drops much more often than with eye ointments. Pressure on the lacrimal punctum for at least a minute after applying eye drops reduces nasolacrimal drainage and therefore decreases systemic absorption from the nasal mucosa.

Also see warnings relating to eye drops and contact lenses.

Eye lotions

These are solutions for the irrigation of the conjunctival sac. They act mechanically to flush out irritants or foreign bodies as a first-aid treatment. Sterile sodium chloride 0.9% solution p. 726 is usually used. Clean water will suffice in an emergency.

Other preparations administered to the eye

Subconjunctival injection may be used to administer anti-infective drugs, mydriatics, or corticosteroids for conditions not responding to topical therapy; intracameral and intravitreal routes can also be used to administer certain drugs, for example antibacterials. These injections should only be used under specialist supervision.

Drugs such as antimicrobials and corticosteroids may be administered systemically to treat susceptible eye conditions.

Ophthalmic Specials

The Royal College of Ophthalmologists and the UK Ophthalmic Pharmacy Group have produced the Ophthalmic Specials Guidance to help prescribers and pharmacists manage and restrict the use of unlicensed eye preparations. 'Specials' should only be prescribed in situations where a licensed product will not be suitable for a child's needs. The Ophthalmic Specials Guidance can be accessed on the Royal College of Ophthalmologists website (www.rcophth.ac.uk). The guidance will be reviewed every six months to ensure the most accurate and up-to-date information is available.

Preservatives and sensitisers

Information on preservatives and substances identified as skin sensitisers is provided under Excipients statements in preparation entries. Very rarely, cases of corneal calcification have been reported with the use of phosphate-containing eye drops in patients with significantly damaged corneas—consult product literature for further information.

Eye preparations: control of microbial contamination

Preparations for the eye should be sterile when issued. Care should be taken to avoid contamination of the contents during use.

Eye drops in multiple-application containers for *domiciliary use* should not be used for more than 4 weeks after first opening (unless otherwise stated by the manufacturer).

Multiple application eye drops for use in *hospital wards* are normally discarded 1 week after first opening—local practice may vary. Individual containers should be provided for each patient. A separate container should be supplied for each eye