

Chapter 13

Skin

CONTENTS

1 Dry and scaling skin disorders	page 765	6 Rosacea and acne	page 805
2 Infections of the skin	772	6.1 Acne	806
2.1 Bacterial skin infections	774	7 Scalp and hair conditions	811
2.2 Fungal skin infections	776	8 Skin cleansers, antiseptics and desloughing agents	812
2.3 Parasitic skin infections	780	8.1 Minor cuts and abrasions	815
2.4 Viral skin infections	781	9 Skin disfigurement	816
3 Inflammatory skin conditions	781	10 Sun protection and photodamage	816
3.1 Eczema and psoriasis	781	11 Superficial soft-tissue injuries and superficial thrombophlebitis	817
4 Perspiration	803	12 Warts and calluses	818
4.1 Hyperhidrosis	803		
5 Pruritus	804		

Skin conditions, management

Topical preparations

When prescribing topical preparations for the treatment of skin conditions in children, the site of application, the condition being treated, and the child's (and carer's) preference for a particular vehicle all need to be taken into consideration.

Vehicles

The British Association of Dermatologists list of preferred unlicensed dermatological preparations (specials) is available at www.bad.org.uk.

The vehicle in topical preparations for the skin affects the degree of hydration, has a mild anti-inflammatory effect, and aids the penetration of the active drug. Therefore, the vehicle, as well as the active drug, should be chosen on the basis of their suitability for the child's skin condition.

Applications are usually viscous solutions, emulsions, or suspensions for application to the skin (including the scalp) or nails.

Collodions are painted on the skin and allowed to dry to leave a flexible film over the site of application.

Creams are emulsions of oil and water and are generally well absorbed into the skin. They may contain an antimicrobial preservative unless the active ingredient or basis is intrinsically bactericidal and fungicidal. Generally, creams are cosmetically more acceptable than ointments because they are less greasy and easier to apply.

Gels consist of active ingredients in suitable hydrophilic or hydrophobic bases; they generally have a high water content. Gels are particularly suitable for application to the face and scalp.

Lotions have a cooling effect and may be preferred to ointments or creams for application over a hairy area. Lotions in alcoholic basis can sting if used on broken skin. *Shake lotions* (such as calamine lotion) contain insoluble powders which leave a deposit on the skin surface.

Ointments are greasy preparations which are normally anhydrous and insoluble in water, and are more occlusive than creams. They are particularly suitable for chronic, dry lesions. The most commonly used ointment bases consist of soft paraffin or a combination of soft, liquid, and hard paraffin. Some ointment bases have both *hydrophilic and lipophilic* properties; they may have occlusive properties on the skin surface, encourage hydration, and also be miscible with water;

they often have a mild anti-inflammatory effect. *Water-soluble ointments* contain macrogols which are freely soluble in water and are therefore readily washed off; they have a limited but useful role where ready removal is desirable.

Pastes are stiff preparations containing a high proportion of finely powdered solids such as zinc oxide and starch suspended in an ointment. They are used for circumscribed lesions such as those which occur in lichen simplex, chronic eczema, or psoriasis. They are less occlusive than ointments and can be used to protect inflamed, lichenified, or excoriated skin.

Dusting powders are used only rarely. They reduce friction between opposing skin surfaces. Dusting powders should not be applied to moist areas because they can cake and abrade the skin. Talc is a lubricant but it does not absorb moisture; it can cause respiratory irritation. Starch is less lubricant but absorbs water.

Dilution

The BP directs that creams and ointments should **not** normally be diluted but that should dilution be necessary care should be taken, in particular, to prevent microbial contamination. The appropriate diluent should be used and heating should be avoided during mixing; excessive dilution may affect the stability of some creams. Diluted creams should normally be used within 2 weeks of preparation.

Suitable quantities for prescribing

Suitable quantities of dermatological preparations to be prescribed for specific areas of the body

Area of body	Creams and Ointments	Lotions
Face	15–30 g	100 ml
Both hands	25–50 g	200 ml
Scalp	50–100 g	200 ml
Both arms or both legs	100–200 g	200 ml
Trunk	400 g	500 ml
Groins and genitalia	15–25 g	100 ml

These amounts are usually suitable for children 12–18 years for twice daily application for 1 week; smaller quantities will be required for children under 12 years. These recommendations **do not apply** to corticosteroid preparations.