

METABOLISM KINETICS

CHARLES W. LOCUSON AND TIMOTHY S. TRACY

University of Minnesota, Minneapolis, Minnesota

Contents

- 1 Introduction
 - 2 Typical Kinetics
 - 2.1 Michaelis–Menten Equation and Its Use in Drug Metabolism
 - 2.2 The Case of Multiple Substrates
 - 2.3 Use of the Michaelis–Menten Equation and Its Parameters
 - 3 Atypical Kinetics Displayed by Drug Metabolizing Enzymes
 - 3.1 Sigmoidal Kinetics with a Single Substrate
 - 3.2 Biphasic Kinetics (Nonasymptotic)
 - 3.3 Hyperbolic (Nonessential) Activation
 - 3.4 Substrate Inhibition
 - 3.5 Explanations for Atypical Kinetics in P450 Enzymes and Their Detection
 - 3.6 Artifactual Sources of Atypical Kinetics
 - 4 Conclusion
- Acknowledgment
References

1 INTRODUCTION

It is common to apply kinetic models to *in vitro* data in an attempt to determine parameters, such as intrinsic clearance, that can be used to make correlations to the *in vivo* situation. These estimations are generally predicated on the data following a hyperbolic profile such that the Michaelis–Menten equation can be applied. This