

Table A.1 Medicines requiring cautionary advisory labels (continued)

Medicine	Ancillary label (number) and/or additional instruction (letter)
Bethanechol	3a or C
Bicalutamide	16, B
Bifonazole	E
Biperiden	1, 9 (long-term regular therapy)
Bisoprolol	9, 12, A (not chewed)
Bosentan	1, 5, 16
Bromazepam	1, 3b, 9
Bromocriptine	5, 12, 16, B
Brompheniramine	1, 13
Budesonide oral capsules	9, A, F
Budesonide oral inhaler with nebuliser solution	14
Bumetanide	16
Buprenorphine	1, 21 (patches)
Bupropion	5, 12†, 16†, A
Buspirone	1, 18
Busulfan	21
Butoconazole	D
Cabergoline	12†, B
Calcitriol	5
Calcium folinate	3b
Candesartan	11, 12†, 16†
Capecitabine	12†, 21, B
Captopril	3b (dose less than 50 mg), 7a (solution), 11, 12, 16†
Carbamazepine	5, 9, 12†, 13, 18, A*, B
Carvedilol	9, 12†, 13, 16
Cefaclor	A*, D
Cefaclor suspension	6, 7a, D
Cefuroxime (oral)	B, D
Celecoxib	10a, 12†
Cephalexin	D
Cephalexin suspension	6, 7a, D
Cetirizine	1, 13
Chloral hydrate	1a, 2, 12
Chlorambucil	6, 12†, 21
Chloramphenicol	7b
Chloroquine	13, B
Chlorpheniramine	1, 13
Chlorpromazine	1, 8, 9 (long-term regular therapy), 16
Chlorthalidone	16, B
Cholecalciferol	5
Ciclesonide	14
Cimetidine	5
Cinacalcet	5, B
Ciprofloxacin (oral)	3b, 4, 8, 12, D
Citalopram	5, 9, 12
Clarithromycin	5, D
Clindamycin	D
Clobazam	1, 9

† Most appropriate during initial treatment or when dosage is increased.

\* Some products have specific indications or specialised formulations or coatings which give rise to instructions different from those applicable generally to the conventional dose form. In cases of doubt concerning specific products with specialised formulations or coatings, reference should be made to the recommendations contained in the manufacturer's information.

Antacids will decrease the absorption of many medicines. Unless specifically indicated to the contrary, the administration of antacids should be separated by a period of about 2 hours from other orally administered medicines whenever possible. Similar separation of the dosing of cholestyramine, colestipol, lanthanum and sevelamer from other medicines is also recommended.