

Table A.1 Medicines requiring cautionary advisory labels (continued)

Medicine	Ancillary label (number) and/or additional instruction (letter)
Disopyramide	9, A
Disulfiram	2, 5
Domperidone	C
Donepezil	12, 16
Dornase alfa	6
Dothiepin	1, 9, 13, 16
Doxepin	1, 9, 13, 16
Doxycycline	4 (delete dairy products), 8, B, D
Doxylamine	1, 13
Droperidol	1
Duloxetine	5, 9, 12, A
Econazole	D*, E (dermal)
Efavirenz	3b, 5, 12
Eformoterol (capsules for inhalation)	22
Emtricitabine	12, A
Enalapril	11, 12†, 16†
Enfuvirtide	12
Entacapone	4 (delete milk, antacids, calcium), 5, 9, 12, 16
Entecavir	3b
Eplerenone	5, 11, 12, 18
Epoetin alfa	6
Epoetin beta	6
Eprosartan	11, 12†, 16†
Erlotinib	3b, 5
Erythromycin	5, A, C, D
Erythromycin ethyl succinate	5, D
Erythromycin ethyl succinate suspension	5, 6, 7a, D
Escitalopram	5, 9, 12
Esomeprazole	A* (can be dispersed in non-carbonated water)
Etanercept	6, 16
Ethacrynic acid	16, B
Ethosuximide	1, 9
Etidronate	4, A, C (2 hours)
Etoposide	18, 21, C
Everolimus	5, 8, 18, A*
Exemestane	B
Exenatide	6, 7b
Ezetimibe	5
Famciclovir	D
Felodipine	9, 12†, 18, A
Fenofibrate	8, A
Fentanyl	1, 21 (patch and lozenge)
Ferrous fumarate	13, B*
Ferrous gluconate	4 (delete dairy products), 13, C
Ferrous sulfate	4 (delete dairy products), 13, A*, C
Flecainide	9, 12, 13
Flucloxacillin	C or 3b, D
Flucloxacillin suspension	C or 3b, 6, 7a, D

† Most appropriate during initial treatment or when dosage is increased.

* Some products have specific indications or specialised formulations or coatings which give rise to instructions different from those applicable generally to the conventional dose form. In cases of doubt concerning specific products with specialised formulations or coatings, reference should be made to the recommendations contained in the manufacturer's information.

Antacids will decrease the absorption of many medicines. Unless specifically indicated to the contrary, the administration of antacids should be separated by a period of about 2 hours from other orally administered medicines whenever possible. Similar separation of the dosing of cholestyramine, colestipol, lanthanum and sevelamer from other medicines is also recommended.