

Table A.1 Medicines requiring cautionary advisory labels (continued)

Medicine	Ancillary label (number) and/or additional instruction (letter)
Procabazine	2, 5, 21, I
Prochlorperazine	1, 16
Proguanil	4 (delete milk, iron and calcium), B
Promethazine	1
Propantheline	9, 12
Propranolol	9, 12†
Quetiapine	1, 9, 12†, 16†
Quinapril	11, 12†, 16†
Quinidine	13, A*
Rabeprazole	A
Raltegravir	12
Ramipril	11, 12†, 16†
Reboxetine	9, 12†, 16
Repaglinide	13
Rifabutin	5, D
Rifampicin	3b, 5, D
Riluzole	12
Risedronate	20 (certain forms), A, C
Risperidone	1, 16
Ritonavir	5, B
Rivastigmine	12, B
Ropinirole	12
Rosiglitazone	10a, A
Rotigotine	1, 9, 12, 13, 16
Roxithromycin	3b, D
Salbutamol	22 (capsules and nebulas), 7b (foil wrapping)
Salcatonin	6
Saquinavir	5, 18, B
Selegiline	5, 12, B
Sertraline	5, 9, 12
Sevelamer	13, A, B
Sibutramine	5, 12
Sildenafil	5, 16
Simvastatin	18
Sirolimus	5, 6 (solution), 7b (solution), 8, 18
Sitaxentan	5
Sodium fusidate	A*, B, D
Somatropin	6
Sorafenib	3b, 5
Sotalol	4 (delete antacids and iron) 9, 12†, C (1–2 hours)
Spironolactone	11, 12†, 16
Strontium ranelate	4
Sucralfate	3b, 5
Sulfadoxine with pyrimethamine	8, B
Sulfasalazine	A*, B
Sulindac	10a, 12†, B
Sulthiame	9
Sumatriptan	12

† Most appropriate during initial treatment or when dosage is increased.

* Some products have specific indications or specialised formulations or coatings which give rise to instructions different from those applicable generally to the conventional dose form. In cases of doubt concerning specific products with specialised formulations or coatings, reference should be made to the recommendations contained in the manufacturer's information.

Antacids will decrease the absorption of many medicines. Unless specifically indicated to the contrary, the administration of antacids should be separated by a period of about 2 hours from other orally administered medicines whenever possible. Similar separation of the dosing of cholestyramine, colestipol, lanthanum and sevelamer from other medicines is also recommended.