

Table A.1 Medicines requiring cautionary advisory labels (continued)

Medicine	Ancillary label (number) and/or additional instruction (letter)
Sunitinib	5, 12
Tacrolimus	3b, 5, 8, 12, 13, 18
Tadalafil	5, 16
Tamoxifen	12†
Tamsulosin	12, 16, A,
Telbivudine	12
Telmisartan	11, 12†, 16†
Temazepam	1 or 1a, 9 (long-term regular therapy)
Temozolomide	21, C (1 hour)
Tenofovir	12†
Terazosin	12†, 16
Terbinafine	D
Teriparatide	6, 7b
Tetrabenazine	1
Thalidomide	1, 16
Theophylline	5, A*, B
Thioguanine	8, 21
Thioridazine	1, 8, 9 (long-term regular therapy), 16
Thyroxine	3b, 4, 6
Tiagabine	1, 9, B
Tiaprofenic acid	10a, 12†, B
Ticlopidine	10b, A, B
Tiludronate	4, A, C (2 hours)
Tinidazole	2, B, D (multiple dosing)
Tiotropium	22 (capsules for inhalation)
Tipranavir	5
Tolterodine	12, 16
Topiramate	1, 9, 12, A*
Tramadol	1, 5, A*
Trandolapril	11, 12†, 16†
Tretinoin (oral)	8, B
Triamterene	11, B
Triazolam	1 or 1a, 9 (long-term regular therapy), 18
Trifluoperazine	1, 8, 9 (long-term regular therapy), 16
Trimeprazine	1, 13
Trimethoprim	D
Trimethoprim Sulfamethoxazole	8, B, D
Trimipramine	1, 9, 13, 16
Tripolidine	1, 13
Tropisetron	12, C (1 hour)
Valaciclovir	D
Valganciclovir	12†, 13, 21, A, B
Valproate	9, 10a, 12†, 13, A*, B
Vancomycin	D
Vardenafil	5, 16
Varenicline	12, 13, A, B
Venlafaxine	5, 9, 12, B
Verapamil	9, 12, 13, 18, A*, B*

† Most appropriate during initial treatment or when dosage is increased.

* Some products have specific indications or specialised formulations or coatings which give rise to instructions different from those applicable generally to the conventional dose form. In cases of doubt concerning specific products with specialised formulations or coatings, reference should be made to the recommendations contained in the manufacturer's information.

Antacids will decrease the absorption of many medicines. Unless specifically indicated to the contrary, the administration of antacids should be separated by a period of about 2 hours from other orally administered medicines whenever possible. Similar separation of the dosing of cholestyramine, colestipol, lanthanum and sevelamer from other medicines is also recommended.