

Table A.1 Medicines requiring cautionary advisory labels

Medicine	Ancillary label (number) and/or additional instruction (letter)
Abacavir	12
Abatacept	6
Acamprosate	2, A, B
Acarbose	B
Acetazolamide	10a, B
Aciclovir (oral)	D
Acitretin	2 (women), 5, 8, B
Adalimumab	6
Adefovir	9
Alendronate	20 (certain dosage forms), A, C
Allopurinol	12†, B
Alprazolam	1, 9
Alprostadil	6 (reconstituted)
Amantadine	9, 12, 16, B
Amiloride	11, 12†, 16
Aminoglutethimide	1†, 16
Aminophylline	5, B
Amiodarone	5, 8, 18
Amisulpride	1, 12, 16
Amitriptyline	1, 9, 13, 16
Amlodipine	9, 12†
Amoxicillin	D
Amoxicillin suspension	6, 7a, D
Amoxicillin with clavulanic acid	13, D, F
Amoxicillin with clavulanic acid suspension	6, 7a, D, F
Amphotericin B	D
Ampicillin	3b, D
Anagrelide	1, 16
Anakinra	6
Anastrozole	12
Aprepitant	5
Aripiprazole	1, 16†
Artemether with lumefantrine	12, A, B
Aspirin	13, A*, B
Atazanavir	5, B
Atenolol	9, 12†
Atomoxetine	5, 12†
Atorvastatin	18
Atovaquone	B, D
Auranofin	B
Azathioprine	8, 21, A*, B
Azithromycin	D (multiple dosing)
Baclofen	1, 9, B
Balsalazide	A, B
Beclomethasone oral inhalers	14
Benzhexol	1, 9 (long-term regular therapy)
Benztropine	1, 9 (long-term regular therapy)
Betahistine	1, 16

† Most appropriate during initial treatment or when dosage is increased.

\* Some products have specific indications or specialised formulations or coatings which give rise to instructions different from those applicable generally to the conventional dose form. In cases of doubt concerning specific products with specialised formulations or coatings, reference should be made to the recommendations contained in the manufacturer's information.

Antacids will decrease the absorption of many medicines. Unless specifically indicated to the contrary, the administration of antacids should be separated by a period of about 2 hours from other orally administered medicines whenever possible. Similar separation of the dosing of cholestyramine, colestipol, lanthanum and sevelamer from other medicines is also recommended.