

Table A.1 Medicines requiring cautionary advisory labels (continued)

Medicine	Ancillary label (number) and/or additional instruction (letter)
Clodronate	3b, (1 hour before or 2 hours after) 4 , A
Clofazimine	8, 12, B
Clomiphene	12
Clomipramine	1, 9, 13, 16
Clonazepam	1, 9
Clonidine	1, 9, 16
Clopidogrel	9, 10a
Clotrimazole	E (dermal)
Clozapine	1, 9, 12, 16
Codeine	1 (greater than 20 mg dose)
Colchicine	5
Cortisone	9, B
Cromoglycate (capsules for inhalation)	22
Cyclophosphamide	8, 21, G
Cyclosporin	5, 8, 18
Cyclosporin (oral solution)	5, 7a, 8, 18
Cyproheptadine	1, 13
Cyproterone	12, B
Dacarbazine	8, 21
Danazol	5, 8
Dantrolene	1, 8
Dapsone	B
Darbepoetin	6
Darunavir	5, B
Dasatinib	4 (antacids), 5, 12, A
Deferasirox	3b, 13
Deferiprone	13
Delavirdine	5
Demeclocycline	3b, 4, 8, D
Desmopressin	5, 13
Desvenlafaxine	9, 12, A
Dexamethasone (oral)	9 (except short courses), B
Dexchlorpheniramine	1, 13, A*
Dextropropoxyphene	1
Diazepam	1 or 1a, 9
Diazoxide	16
Diclofenac	10a, 12†, A*, B
Dicloxacillin	3a, D
Didanosine	3b, A*
Didanosine suspension	3b, 6, 7a
Digoxin	5
Dihydrocodeine	1 (greater than 20 mg dose)
Dihydroergotamine	5, 18
Diltiazem	9, 12†, A*
Dimenhydrinate	1, 13
Diphenhydramine	1, 13
Diphenoxylate	1, 13
Dipyridamole	16, A*

† Most appropriate during initial treatment or when dosage is increased.

* Some products have specific indications or specialised formulations or coatings which give rise to instructions different from those applicable generally to the conventional dose form. In cases of doubt concerning specific products with specialised formulations or coatings, reference should be made to the recommendations contained in the manufacturer's information.

Antacids will decrease the absorption of many medicines. Unless specifically indicated to the contrary, the administration of antacids should be separated by a period of about 2 hours from other orally administered medicines whenever possible. Similar separation of the dosing of cholestyramine, colestipol, lanthanum and sevelamer from other medicines is also recommended.