

- To avoid hazardous activities until product response is determined
- That orthostatic hypotension occurs often; to rise from sitting or lying position gradually
- To avoid hot tubs, hot showers, tub baths because hypotension may occur
- To avoid abrupt withdrawal of this product because EPS may result; product should be withdrawn slowly
- To avoid OTC preparations (cough, hay fever, cold) unless approved by prescriber because serious product interactions may occur; to avoid use with alcohol, CNS depressants, opioids because increased drowsiness may occur
- **Suicidal thoughts, behaviors: report immediately suicidal thoughts/behaviors**
- **That in hot weather, heat stroke may occur; to take extra precautions to stay cool**
- To take PO without regard to food
- To notify prescriber if pregnancy is planned or suspected; not to breastfeed

TREATMENT OF OVERDOSE: Lavage if orally ingested; provide airway; do not induce vomiting or use EPINEPHrine

⚠ HIGH ALERT

RARELY USED

olaparib

(oh-lap'a-rib)

Lynparza

Func. class.: Antineoplastic-PARP

USES: Treatment of deleterious or suspected deleterious germline BRCA-mutated advanced ovarian cancer in patients who have not responded successfully to ≥ 3 prior courses of chemotherapy, as monotherapy

CONTRAINDICATIONS: Hypersensitivity

DOSAGE AND ROUTES

- **Adult female: PO** 400 mg bid, (tablets) 300 mg bid (capsules) until disease

progression or unacceptable toxicity. Avoid use of concomitant strong and moderate CYP3A4 inhibitors if possible

⚠ HIGH ALERT

RARELY USED

olaratumab

(oh-lar-at'ue-mab)

Lartruvo 

Func. class.: Antineoplastic, monoclonal antibody

USES: The treatment of soft-tissue sarcoma not amenable to curative treatment with radiotherapy or surgery, in combination with DOXOrubicin


CONTRAINDICATIONS: Hypersensitivity, pregnancy

DOSAGE AND ROUTES

- **Adult: IV INFUSION** 15 mg/kg over 60 min on days 1 and 8 repeated q21days until disease progression or unacceptable toxicity in combination with doxorubicin 75 mg/m² **IV** on day 1 repeated q21days for up to 8 cycles

olmesartan (Rx)

(ol-meh-sar'tan)

Benicar, Olmetec 

Func. class.: Antihypertensive

Chem. class.: Angiotensin II receptor (type AT₁) antagonist

Do not confuse:

Benicar/Mevacor

ACTION: Blocks the vasoconstrictor and aldosterone-secreting effects of angiotensin II; selectively blocks the binding of angiotensin II to the AT₁ receptor found in tissues

USES: Hypertension, alone or in combination with other antihypertensives

CONTRAINDICATIONS: Hypersensitivity