

- That if a dose is missed, skip missed dose and administer the next dose the following day in the morning on wakening
- To report to healthcare provider if inability to stay awake continues or if you develop anxiety, agitation, irritability problems sleeping
- **Pregnancy/breastfeeding:** identify if pregnancy is planned to suspected, encourage to enroll in the pregnancy registry if she becomes pregnant; to enroll or obtain information from the registry, online at www.SunosiPregnancyRegistry.com or by calling 1-877-283-6220; present in breast milk

tenapanor

(ten-a' pa- nor)

Ibsrela

Func. class.: IBS agent

USES: Irritable bowel syndrome with constipation in adults

CONTRAINDICATIONS

Hypersensitivity, GI obstruction

DOSAGE AND ROUTES

- **Adult:** **PO** 50 mg bid immediately before breakfast or the first meal of the day and immediately before dinner

ubrogepant

Ubrelyv

Func. class.: Calcitonin gene–related peptide (CGRP) receptor antagonist

USES: Acute migraine

DOSAGE AND ROUTES

- **Adult:** **PO** 50 or 100 mg; If needed, may take second dose at least 2 hr after initial dose; max: 200 mg/24 hr

CONTRAINDICATIONS

Hypersensitivity, strong CYP3A4 inhibitors

upadacitinib (Rx)

(Ue-pad' a-sye' ti-nib)

Rinvoq

Func. class.: Immunomodulating agent*Chem. class.:* Janus associated kinase (JAK) inhibitor

ACTION: Oral JAK inhibitor; Janus kinase, a tyrosine kinase enzyme that transmit signals arising from cytokine or growth factor–receptor interactions on the cellular membrane to influence cellular processes of immune cell function and hematopoiesis

USES: Treatment of rheumatoid arthritis

CONTRAINDICATIONS

Hypersensitivity

Precautions: AIDS, anemia, breastfeeding, children, contraception requirements, corticosteroid use, diverticulitis, geriatrics, GI perforation, hepatic disease, hepatitis B exacerbation, herpes, HIV, neutropenia, pregnancy, pregnancy testing, vaccination

Black Box Warning: Infection, new primary malignancy, thrombosis

DOSAGE AND ROUTES

- **Adults:** **PO** 15 mg daily with or without methotrexate or other nonbiologic DMARDs

Available forms: Extended release tablet 15 mg

Administer:

- Give with or without food
- Swallow whole; do not split, crush or chew

NURSING CONSIDERATIONS**Assess:**

- **RA:** pain, stiffness, ROM, swelling of joints before, during treatment
- Infections (fever, flulike symptoms, dyspnea, change in urination, redness/