

TREATMENT OF OVERDOSE:

Lavage; provide an airway; do not induce vomiting

RARELY USED**cobicistat/darunavir/
emtricitabine/tenofovir**

Symtuza

Func. class.: Antiviral-HIV protease inhibitor/nucleotide reverse transcriptase inhibitor (NRTI) combination

USES: Human immunodeficiency virus (HIV) infection in treatment-naïve adults/virologically suppressed adults on an antiretroviral regimen ≥ 6 mo and who are without resistance-associated substitutions to darunavir or tenofovir

CONTRAINDICATIONS: Hypersensitivity

Black Box Warning: Hepatitis B exacerbation

DOSAGE AND ROUTES

• **Adult/adolescent:** **PO** 1 tablet (800 mg darunavir; 150 mg cobicistat; 200 mg emtricitabine; 10 mg tenofovir alafenamide) daily with food

⚠ HIGH ALERT**RARELY USED****cobimetinib**

(koe-bi-me' ti-nib)

Cotellic

Func. class.: Antineoplastic

USES: Orphan drug. For the treatment of unresectable or metastatic melanoma in patients with a BRAF V600E or V600K mutation, in combination with vemurafenib

CONTRAINDICATIONS: Hypersensitivity

DOSAGE AND ROUTES

• **Adult:** **PO** 60 mg (three 20-mg tablets) q day \times 21 days, in combination with vemurafenib 960 mg bid \times 28 days; repeat cycle q28days until disease progression or unacceptable toxicity

**codeine (Rx)**

(koe'deen)

Func. class.: Opiate analgesic, antitussive

Chem. class.: Opiate, phenathrene derivative

Controlled Substance Schedule II, III, IV, V (depends on content)

Do not confuse:

codeine/Lodine/iodine

ACTION: Depresses pain impulse transmission at the spinal cord level by interacting with opioid receptors; decreases cough reflex, GI motility

USES: Mild to moderate pain

Unlabeled uses: Diarrhea

CONTRAINDICATIONS: Breast-feeding, hypersensitivity to opiates, respiratory depression, increased intracranial pressure, seizure disorders, severe respiratory disorders

Black Box Warning: Children recovering from tonsillectomy/adenoidectomy who are ultrarapid metabolizers

Precautions: Pregnancy, geriatric patients, cardiac dysrhythmias, prostatic hypertrophy, bowel impaction

DOSAGE AND ROUTES**Pain**

• **Adult:** **PO** 15-60 mg q4hr prn

• **Child 6-17 yr:** **PO** 3 mg/kg/day in divided doses q4hr prn