


naphazoline
(ophthalmic)

(na-faz'oh-leen)

Advanced Eye Relief, Redness Maximum Relief, Ak-Con, All Clear, Clear Eyes, Naphcon Forte , VasoClear*Func. class.:* Ophthalmic vasoconstrictor*Chem. class.:* Sympathomimetic**ACTION:** Acts on the blood vessels in the eye to produce vasoconstriction**USES:** Ocular congestion, irritation, itching of the eye**CONTRAINDICATIONS:** Hypersensitivity, acute angle-closure glaucoma, 0.1% solution in children/infants**Precautions:** Hyperthyroidism, diabetes mellitus, hypertension, cardiac conditions**DOSAGE AND ROUTES**

- **Adult:** **OPHTH** instill 1-2 drops in affected eye in the conjunctival sac every 3-4 hr, up to qid

Available forms: ophthalmic solution 0.012%, 0.1%**Administer:**

- Store at room temperature; keep tightly closed

SIDE EFFECTS**CNS:** Headache**EENT:** Blurred vision, irritation, photophobia, dilation, stinging, elevated IOP, keratitis**PHARMACOKINETICS**

Onset 10 min, duration up to 6 hr

INTERACTIONS**Increase:** Systemic effects— β -blockers

- Do not use within 14 days of MAOIs

NURSING CONSIDERATIONS**Assess:**

- Ocular itching, congestion, irritation: should show improvement quickly; avoid using more than 3 consecutive days; long-term use or exceeding dosage can

lead to rebound congestion; report eye pain, blurred vision

Evaluate:

- Decreasing ocular itching, congestion, irritation

Teach patient/family:

- Method for instilling drops
- **To notify prescriber of eye pain, blurred vision**
- That ocular itching, congestion, irritation should show improvement quickly
- To avoid using longer than 3 days; that long-term use or exceeding dosage can lead to rebound congestion
- To wait for at least 15 min before wearing contact lenses

naproxen (Rx, OTC)Aleve, Anaprox, Anaprox DS, EC-Naprosyn, Maxidol , Naprelan, Naprosyn*Func. class.:* Nonsteroidal antiinflammatory, nonopioid analgesic*Chem. class.:* Propionic acid derivative**ACTION:** Inhibits COX-1, COX-2 by blocking arachidonate; analgesic, antiinflammatory, antipyretic**USES:** Osteoarthritis; rheumatoid, gouty arthritis; primary dysmenorrhea; ankylosing spondylitis, bursitis, tendinitis, myalgia, dental pain, juvenile rheumatoid arthritis**Unlabeled uses:** Bone pain, migraine/migraine prophylaxis, heterotropic ossification**CONTRAINDICATIONS:** Pregnancy 3rd trimester; hypersensitivity to NSAIDs, salicylates; perioperative pain in CABG surgery; MI; stroke**Precautions:** Pregnancy, breastfeeding, children <2 yr, geriatric patients, bleeding disorders, GI disorders, cardiac disorders, hypersensitivity to other antiinflammatory agents, CCr <30 mL/min, asthma, renal failure, hepatic disease

N