

370 deoxycholic acid

should be given to prevent osteonecrosis of the jaw

- **Infection:** do not start treatment in patients with active infections; infections should be resolved first
- **Pregnancy:** identify whether pregnancy is planned or suspected; women who become pregnant should enroll in Amgen's Pregnancy Surveillance Program (1-800-772-6436); do not use in pregnancy, breastfeeding

Evaluate:

- Therapeutic response: increased/maintained bone density, decreased calcium levels

Teach patient/family:

- To report hypercalcemic relapse: nausea, vomiting, bone pain, thirst
- To notify prescriber immediately if rash, infection, cramps, twitching occur
- To continue with dietary recommendations, including additional Ca 1000 mg/day and vit D \geq 400 units (Prolia product labeling)
- That product must be continued or fractures may occur
- To use acetaminophen before and for 72 hr after inj to lessen bone pain
- About the purpose of this product and its expected results
- To avoid OTC, Rx medications and herbs and supplements unless approved by prescriber
- To exercise regularly, stop smoking, and avoid alcohol to maintain bone health
- To inform all health care providers of product use; to avoid dental procedures/surgery if possible; to practice good oral hygiene
- That lab tests and follow-up exams will be required
- **Pregnancy/breastfeeding:** not to use during pregnancy and breastfeeding; to notify prescriber if pregnancy is planned, suspected; to use contraception during and for (Xgeva) 5 mo after completion of therapy; that males

should use contraception if partner is pregnant

RARELY USED

deoxycholic acid

(dee-ox'-i-koe'-lik as'-id)

Kybella 

Func. class.: Lypolytic

USES: For the improvement in the appearance of moderate to severe convexity or fullness associated with submental fat

CONTRAINDICATIONS: Infection

DOSAGE AND ROUTES

- **Adult:** **SUBCUT** 0.2 mL per injection site. A single treatment session consists of up to a maximum of 50 injections (10 mL total dose), with each injection spaced 1 cm apart. Up to 6 single treatments may be given at intervals of no less than 1 mo. Injections are made into the subcutaneous fat of the submental region, between the dermis and the platysma. Do not inject into the platysma

desipramine (Rx)

(dess-ip'ra-meen)

Norpramin

Func. class.: Antidepressant, tricyclic

Chem. class.: Dibenzazepine, secondary amine

Do not confuse:

desipramine/disopyramide

ACTION: Blocks reuptake of norepinephrine, serotonin into nerve endings, thereby increasing action of norepinephrine, serotonin in nerve cells

USES: Depression

Unlabeled uses: Chronic pain, insomnia, anxiety

CONTRAINDICATIONS: Hypersensitivity to tricyclics, carbamazepine, closed-angle glaucoma, acute MI, MAOIs