
2 Mathematical Models in Percutaneous Absorption

Yuri G. Anissimov

Griffith University, Brisbane, Australia

Griffith University, Gold Coast, Australia

Joshua J. Calcutt

Griffith University, Brisbane, Australia

Michael S. Roberts

The University of Queensland, Princess Alexandra

Hospital, Woolloongabba, Australia

University of South Australia, Adelaide, Australia

CONTENTS

2.1	In Vitro Skin Diffusion Models in Percutaneous Absorption	12
2.1.1	In Vitro Skin Permeability Studies with a Constant Donor Concentration and Sink Receptor Conditions	12
2.1.2	Amount and Flux–Time Profiles on Removing the Donor Phase After Reaching the Steady State for Conditions Described in Section 2.1.2.A	16
2.1.3	In Vitro Permeability Studies with a Constant Donor Concentration and Finite Receptor Volume	18
2.1.4	In Vitro Permeability Studies with a Constant Donor Concentration or Defined Input Flux and Finite Clearance of Solute from the Epidermis	20
2.1.5	In Vitro Skin Permeability Studies with Finite Donor Volume and Receptor Sink Conditions	21
2.1.6	In Vitro Permeability Studies with a Finite Donor Volume and a Finite Clearance from the Epidermis into the Receptor	24
2.1.7	In Vitro Skin Permeability Studies with Diffusion Limited Finite Donor and Sink Receptor Conditions	24
2.1.8	In Vitro Permeability Studies with Two-Layer Diffusion Limitations in Transport	26
2.1.9	Desorption	27
2.1.10	Modeling the Corneocytes and Lipids Explicitly	28
2.1.11	Slow Binding in SC	29
2.1.12	SC Heterogeneity	29
2.2	Release Profiles from Topical Products	30
2.2.1	Diffusion-Controlled Release	30
2.2.2	Release of a Suspended Drug by Diffusion	30
2.3	Compartmental Models as an Alternative to Diffusion Models in Percutaneous Absorption	31