


In addition to the saponin, α -spinasteryl glucoside and 7,3',4'-trihydroxyflavone were found in the stem bark of *A. julibrissin* [5]. From the lower of *A. julibrissin* cyanidin-3-glucoside (9-6) was identified [6].



Cyanidin-3- β -D-glucopyranoside (9-6)

9.3 Pharmacology

The saponin fraction of the stem bark of *A. julibrissin*, from which machaerinic acid methyl ester and acacid acid lactone were isolated, had a strong uterotonic activity [2].

References

1. Wang CS (1982) Chemical identification of some traditional Chinese drugs containing saponins. *Bull Chin Mat Med* 7:13–14
2. You YH, Woo WS, Choi JS, Kang SS (1982) Isolation of a new saponin from *Albizzia julibrissin*. *Arch Pharmacol Res* 5:33–38 (CA 98:122814n)
3. Kang SS, Woo WS (1983) Saponin from *Albizzia julibrissin*. *Arch Pharmacol Res* 6:25–28 (CA 99:102292h)
4. Woo WS, Kang SS (1984) Isolation of a new monoterpene conjugated triterpenoid from the stem bark of *Albizzia julibrissin*. *J Nat Prod* 47:547–549
5. Chamsuksai P, Choi JS, Woo WS (1981) 3',4',7-Trihydroxyflavone in *Albizzia julibrissin*. *Arch Pharmacol Res* 4:1219–1231 (CA 96:196564m)
6. Ishikura N, Ito S, Shibata M (1978) Paper chromatographic survey of anthocyanins in Leguminosae. III. Identification and distribution of pattern of anthocyanins in twenty-two legumes. *Bot Mag* 91:25–30 (CA 89:103708d)