

NEOSPORIN CREAM (neomycin—topical + polymyxin—topical + bacitracin—topical) Apply one to three times per day. [OTC Trade only: neomycin 3.5 mg/g + polymyxin 10,000 units/g; 15 g and unit dose 0.94 g.] ▶K ♀D? \$

NEOSPORIN OINTMENT (bacitracin—topical + neomycin—topical + polymyxin—topical) Apply one to three times per day. [OTC Generic/Trade: bacitracin 400 units/g + neomycin 3.5 mg/g + polymyxin 5000 units/g 15, 30 g and “to go” 0.9 g packets.] ▶K ♀D? \$

OZENOXACIN (*Xepi*) **Impetigo**: Apply to affected area twice daily for 5 days. [Rx/Trade: cream 1%, 10, 30, 45 g.] ▶Not absorbed ♀ +/+ + D + \$\$\$\$\$

POLYSPORIN (bacitracin—topical + polymyxin—topical, **Polytopic*) Apply one to three times per day. [OTC Trade only: Oint 15, 30 g and unit dose 0.9 g. Powder 10 g.] ▶K ♀D? \$

RETAPAMULIN (*Altanax*) **Impetigo**: Apply thin layer two times per day for 5 days. [Trade only: Oint 1% 15, 30 g.] ▶Not absorbed ♀B D? \$\$\$\$\$

SILVER SULFADIAZINE (*Silvadene*, **Flamazine*) Apply one to two times per day. [Generic/Trade: Cream 1% 20, 50, 85, 400, 1000 g.] ▶LK ♂♀B D— \$ \$

Antifungals (Topical)

BUTENAFINE (*Lotrimin Ultra*, *Mentax*) **Treatment of tinea pedis**: Apply daily for 4 weeks or two times per day for 7 days. **Tinea corporis, tinea versicolor, or tinea cruris**: Apply daily for 2 weeks. [Rx Trade only: Cream 1% 15, 30 g (*Mentax*). OTC Generic/Trade: Cream 1% 12, 24 g (*Lotrimin Ultra*).] ▶L ♀B D? \$

CICLOPIROX (*Loprox*, *Loprox TS*, *Penlac*, **Stieprox shampoo*) **Tinea pedis, tinea cruris, tinea corporis, tinea versicolor, and candidiasis (cream, lotion)**: Apply two times per day. **Fungal nail infection (Penlac)**: Apply daily to affected nails; apply over previous coat; remove with alcohol every 7 days. **Seborrheic dermatitis (Loprox shampoo)**: Shampoo two times per week for 4 weeks. [Generic/Trade: Shampoo (*Loprox*) 1% 120 mL. Nail soln (*Penlac*) 8% 6.6 mL. Generic only: Gel 0.77% 30, 45, 100 g. Cream (*Loprox*) 0.77% 15, 30, 90 g. Lotion (*Loprox TS*) 0.77% 30, 60 mL.] ▶K ♀B D? \$\$\$\$

CLOTRIMAZOLE—TOPICAL (*Lotrimin AF*, *Mycelex*, **Canesten*, *Clotrimaderm*) **Treatment of tinea pedis, tinea cruris, tinea corporis, tinea versicolor, and cutaneous candidiasis**: Apply two times per day. [Note that different Lotrimin brand name products may contain clotrimazole, miconazole, or butenafine. Rx Generic only: Cream 1% 15, 30, 45 g. Soln 1% 10, 30 mL. OTC Generic/Trade (*Lotrimin AF*): Cream 1% 12, 24 g.] ▶L ♀B D? \$

ECONAZOLE (*Ecoza*) **Tinea pedis, tinea cruris, tinea corporis, tinea versicolor**: Apply daily. **Cutaneous candidiasis**: Apply two times per day. [Generic only: Cream 1% 15, 30, 85 g. Trade only: Foam 1% 70 g (*Ecoza*—\$\$\$\$).] ▶Not absorbed ♀C D? \$\$\$

FINACONAZOLE (*Jublia*) **Onychomycosis of toenail**: Apply once daily to affected toenail for 48 weeks. [Trade: Soln 1% 4, 8 mL brush applicator.] ▶Minimal absorption ♀C D? \$\$\$\$\$