

**GLYCOPYRRONIUM (Qbrexza)** **Primary axillary hyperhidrosis:** Apply to each axillae once daily using the same cloth. [Single-use cloth with 2.4% glycopyrronium.] ▶? ♀?!? ♂?

**HYALURONIC ACID (Bionect, Restylane, Perlane)** **Moderate to severe facial wrinkles:** Inject into wrinkle/fold (Restylane, Perlane). **Protection of dermal ulcers:** Apply gel/cream/spray two or three times per day (Bionect). [OTC Trade only: Cream 2% 15, 30 g. Rx Generic/Trade: Soln 3% 30 mL. Gel 4% 30 g. Cream 4% 15, 30, 60 g. Injectable gel 2%.] ▶? ♀? ♂? \$\$\$

**HYDROGEN PEROXIDE (Eskata)** **Raised seborrheic keratosis lesions:** Applied by health care practitioner only. Apply four times, at 1-min intervals. [Rx/Trade: soln, 4% in a single-use applicator.] ▶Not absorbed ♀+!/+! ♂+ \$\$\$\$\$

**HYDROQUINONE (Eldopaque, Eldoquin, Eldoquin Forte, EpiQuin Micro, Esoterica, Glyquin, Lustra, Melanex, Solaquin, Claripel, \*Ultraquin)** **Hyperpigmentation:** Apply two times per day. [OTC Trade only: Cream 2% 15, 30 g. Rx Generic/Trade: Soln 3% 30 mL. Gel 4% 30 g. Cream 4% 15, 30, 60 g.] ▶? ♀? ♂? \$

**IVERMECTIN—TOPICAL (Sklice, Soolantra)** **Lice (Sklice):** Apply lotion to dry hair and scalp. Rinse after 10 min. Single application only. **Rosacea (Soolantra):** Apply cream to affected area once daily. [Trade only: Lotion (Sklice) 0.5%, 120 mL. Cream (Soolantra) 1% 30, 45, 60 g.] ▶Minimal absorption ♀? ♂? \$\$\$\$\$

**LACTIC ACID (Lac-Hydrin, AmLactin, \*Dermalac)** Apply two times per day. [Trade only: Lotion 12% 150, 360 mL. OTC: Cream 12% 140, 385 g. AmLactin AP is lactic acid (12%) with pramoxine (1%).] ▶? ♀? ♂? \$\$\$

**LIDOCAINE—TOPICAL (Ztlido, Xylocaine, Lidoderm, Numby Stuff, LMX, Zingo, \*Maxilene)** Apply prn. Dose varies with anesthetic procedure, degree of anesthesia required, and individual patient response. **Postherpetic neuralgia (patch):** Apply up to 3 patches to affected area at once for up to 12 h within a 24-h period. (Lidoderm, Ztlido) Apply 30 min prior to painful procedure (ELA-Max 4%). **Discomfort with anorectal disorders:** Apply prn (ELA-Max 5%). **Intradermal powder injection for venipuncture/IV cannulation, 3 to 18 yo (Zingo):** 0.5 mg to site 1 to 10 min prior. [For membranes of mouth and pharynx: Spray 10%, Oint 5%, Liquid 5%, Soln 2%, 4%, Dental patch. For urethral use: Jelly 2%. Patch (\$\$\$\$\$): Lidoderm 5%, Ztlido 1.8%. Intradermal powder injection system: 0.5 mg (Zingo). OTC Trade only: Liposomal lidocaine 4% (ELA-Max). Patches 4%, various brands.] ▶LK ♀B ♂+ varies by therapy

**MINOXIDIL—TOPICAL (Rogaine, Women's Rogaine, Rogaine Extra Strength, Minoxidil for Men)** **Androgenetic alopecia in men or women:** 1 mL to dry scalp two times per day (2% soln) or once daily (5% soln). [OTC Generic/Trade: Soln 2% 60 mL (Rogaine, Women's Rogaine). Soln 5% 60 mL (Rogaine Extra Strength, Theroxidil Extra Strength—for men only). Foam 5% 60 g (Rogaine Extra Strength).] ▶K ♀C ♂—\$

**OATMEAL (Aveeno)** **Pruritus from poison ivy/oak, varicella:** Apply lotion four times per day prn. Also bath packets for tub. [OTC Generic/Trade: Lotion. Bath packets.] ▶Not absorbed ♀? ♂? \$

**OXYMETAZOLINE—TOPICAL (Rhofade)** **Persistent facial erythema associated with rosacea:** Apply pea-sized amount to entire face once daily. Avoid eyes and lips. [Trade: Rx, cream 1%, 30 g, 60 g tubes and pump.] ▶Minimal absorption ♀?!?!? ♂? \$\$\$\$\$