

OBETICHOIC ACID (*Ocaliva*) **Primary biliary cholangitis, with ursodiol if no response to ursodiol for at least one year or without ursodiol in those intolerant to ursodiol:** 5 mg PO once daily. Can be increased to 10 mg once daily in 3 months, if inadequate response. [Trade only: Tabs 5, 10 mg.] ▶L, Bile ♀?/?/? ♀? \$\$\$\$\$

OCETROTIDE (*Sandostatin, Sandostatin LAR*) **Variceal bleeding:** Bolus 25 to 50 mcg IV followed by infusion 25 to 50 mcg/h. **AIDS diarrhea:** 25 to 250 mcg SC three times per day. [Generic: Injection vials 0.05, 0.1, 0.2, 0.5, 1 mg. Trade only: Long-acting injectable susp (Sandostatin LAR) 10, 20, 30 mg.] ▶LK Ⓞ ♀B ♀? \$\$\$\$\$

ORLISTAT (*Alli, Xenical*) **Weight loss:** 60 to 120 mg PO three times per day with meals. [OTC Trade only (Alli): Caps 60 mg. Rx Trade only (Xenical): Caps 120 mg.] ▶gut ♀X ♀? \$\$\$

PANCREATIN (*Creon, Ku-Zyme*) 8000 to 24,000 units lipase (1 to 2 tabs/caps) PO with meals and snacks. [Tabs, Caps with varying amounts of lipase, amylase, and protease.] ▶gut ♀C ♀? \$\$\$

PANCRELIPASE (*Creon, Pancreaze, Pancrecarb, Cotazym, Ku-Zyme HP, Ultresa, Viokace, Zenpep*) Varies by wt. Initial infant dose 2000 to 4000 lipase units PO per 120 mL formula or breastmilk. 12 mo or older to younger than 4 yo: 1000 lipase units/kg PO. 4 yo or older: 500 lipase units/kg per meal PO, max 2500 lipase units/kg per meal. [Tabs, Caps, Powder with varying amounts of lipase, amylase, and protease.] ▶gut ♀C ♀? \$\$\$

•**PINAVERIUM** (•*Dicetel*) Canada only. 50 to 100 mg PO three times per day. [Trade only: Tabs 50, 100 mg.] ▶? ♀C ▶—\$\$\$

PLECANATIDE (*Trulanc*) **Chronic idiopathic constipation, IBS with constipation:** 3 mg PO once daily. Avoid in children 17 yo or younger. [Trade only: tab 3 mg.] ▶gut ♀?/?/? ♀? \$\$\$\$\$ ■

QSYMIA (phentermine + topiramate) **Obesity (BMI 30 or greater) or overweight (BMI 27 or greater with weight-related comorbidities):** 3.75 mg/23 mg PO once daily for 14 days, then increase to 7.5 mg/46 mg PO once daily. Max dose 15 mg/92 mg PO once daily. Taper over 1 week when discontinuing. [Trade: Tabs 3.75/23, 7.5/46, 11.25/69, 15/92 mg (phentermine/topiramate).] ▶KL Ⓞ ♀X ▶—ⓄIV \$\$\$\$\$

RECTIV (*nitroglycerin—rectal*) **Painful chronic anal fissures:** Apply 1 inch intrarectally q 12 for up to 3 weeks. [Trade only: Oint 0.4% 30 g.] ▶L ♀C ♀? \$\$\$\$\$

SECRETIN (*ChiRhoStim*) 0.2 to 0.4 mcg/kg IV over 1 min. ▶Serum ♀?/?/? ♀? \$\$\$\$\$

TEDUGLUTIDE (*Gattex, •Revestive*) **Short bowel syndrome patients receiving IV TPN:** 0.05 mg/kg (max 3.8 mg) SC daily. [Trade only: 5 mg/vial, powder for reconstitution.] ▶endogenous Ⓞ ♀B ♀? \$\$\$\$\$

URSODIOL (*Actigall, URSO, URSO Forte*) **Gallstone solution** (Actigall): 8 to 10 mg/kg/day PO divided two to three times per day. **Prevention of gallstones associated with rapid wt loss** (Actigall): 300 mg PO two times per day. **Primary biliary cirrhosis** (URSO): 13 to 15 mg/kg/day PO divided in 2 to 4 doses. [Generic/Trade: Caps 300 mg, Tabs 250, 500 mg.] ▶Bile ♀B ♀? \$\$\$\$\$