

**GUSELKUMAB (Tremfya)** 100 mg SC at week 0 and 4, then every 8 weeks thereafter. [Trade: Inj soln: 100 mg/1 mL prefilled syringe.] ▶endogenous ♀?/?/? ♯? \$\$\$\$\$

**IXEKIZUMAB (Taltz)** Moderate to severe plaque psoriasis: 160 mg SC at week 0, then 80 mg SC at weeks 2, 4, 6, 8, 10, and 12, then 80 mg SC every 4 weeks. Psoriatic arthritis: 160 mg SC at week 0, then 80 mg SC every 4 weeks. For psoriatic arthritis patients with moderate-to-severe plaque psoriasis, use the dosing regimen for plaque psoriasis. [Trade only: prefilled autoinjector 80 mg, prefilled syringe 80 mg.] ▶proteolysis ♀?/?/? ♯? \$\$\$\$\$

**TACLONEX (calcipotriene + betamethasone—topical)** Apply daily for up to 4 weeks. [Calcipotriene 0.005% + betamethasone dipropionate 0.064%. Generic/Trade: Oint 60, 100 g. Trade only: Topical susp (Taclonex) 60, 120 g.] ▶L ♀ ♯? \$\$\$\$\$

**TILDRAKIZUMAB (Ilumya)** Moderate to severe plaque psoriasis: 100 mg SC at weeks 0, 4, and every 12 weeks thereafter. [Trade only: 100 mg prefilled syringe.] ▶endogenous ♀?/?/? ♯? \$\$\$\$\$

**USTEKINUMAB (Stelara)** Severe plaque psoriasis, active psoriatic arthritis, wt less than or equal to 100 kg, 12 yo or older: 45 mg SC initially and again 4 weeks later, followed by 45 mg SC q 12 weeks. For wt greater than 100 kg and psoriatic arthritis with moderate to severe plaque psoriasis, 12 yo or older: 90 mg SC initially and again 4 weeks later, followed by 90 mg SC q 12 weeks. Moderate to severe active Crohn's Disease: Up to 55 kg: 260 mg as a single IV infusion, 56 kg to 85 kg: 390 mg as a single IV infusion, 85 kg or greater: 520 mg as a single IV infusion. Maintenance in moderate to severe Crohn's Disease: 90 mg SC starting 8 weeks after IV infusion, then every 8 weeks. [Trade only: 45 and 90 mg prefilled syringe and vial. Single-dose vial, 130 mg/26 mL solution for infusion.] ▶L ♀B ♯? \$\$\$\$\$

### Antivirals (Topical)

**ACYCLOVIR—TOPICAL (Zovirax)** Herpes genitalis: Apply ointment q 3 h (6 times per day) for 7 days. Recurrent herpes labialis: Apply cream 5 times per day for 4 days. [Generic/Trade: Oint 5% 5, 15, 30 g. Trade only: Cream 5% 5 g.] ▶K ♀C ♯? \$\$\$\$\$

**DOCOSANOL (Abreva)** Oral-facial herpes (cold sores): Apply 5 times per day until healed. [OTC Trade only: Cream 10% 2 g.] ▶Not absorbed ♀B ♯? \$

**IMIQUIMOD (Aldara, Zyclara, +Vyloma)** Genital/perianal warts: Apply once daily for up to 8 weeks. Wash off after 8 h. Non-hyperkeratotic, non-hypertrophic actinic keratoses on face/scalp in immunocompetent adults: Apply two times per week overnight for 16 weeks (Aldara) or once daily for two 2-week periods separated by a 2-week break (Zyclara). Wash off after 8 h. Primary superficial basal cell carcinoma: Apply 5 times a week for 6 weeks (Aldara). Wash off after 8 h. [Generic/Trade: Cream 5% (Aldara) single-use packets. Trade only: Cream 3.75% (Zyclara-\$\$\$\$\$) single-use packets, box of 28 and 7.5 g pump. Cream 2.5% 7.5 g pump.] ▶Not absorbed ♀C ♯? \$\$\$

**PENCICLOVIR (Denavir)** Herpes labialis (cold sores): Apply cream q 2 h while awake for 4 days. [Trade only: Cream 1% tube 5 g.] ▶Not absorbed ♀B ♯? \$\$\$\$\$