

**Glaucoma Agents—Combinations and Other**

**COMBIGAN** (brimonidine + timolol—ophthalmic) 1 gtt in each affected eye two times per day. Contraindicated in children younger than 2 yo. [Trade only: Soln brimonidine 0.2% + timolol 0.5% (5, 10, 15 mL).] ▶LK ♀C ▶—\$\$\$\$

**COSOPT** (dorzolamide + timolol—ophthalmic) 1 gtt in each affected eye two times per day. [Generic/Trade: Soln dorzolamide 2% + timolol 0.5% (10 mL). Trade only: Soln, preservative-free dorzolamide 2% + timolol 0.5% (60 single-use containers).] ▶LK ♂♀D ▶—\$\$\$

**NETARSUDIL** (*Rhopressa*) **Reduction of IOP in open-angle glaucoma or ocular HTN:** One drop in affected eye(s) every evening. [Rx/Trade: soln 0.02%, 2.5 mL.] ▶Not absorbed ♀?/?/? ▶? \$\$\$\$\$

**SIMBRINZA** (brinzolamide + brimonidine) 1 gtt in each affected eye three times per day. [Trade only: Brinzolamide 1% and brimonidine 0.2% 8 mL.] ▶LK ♂♀C ▶? \$\$\$\$

**Glaucoma Agents—Miotics**

**CARBACHOL** (*Isopto Carbachol, Miostat*) [Trade only: Soln (Isopto Carbachol) 1.5, 3% (15 mL). Intraocular soln (Miostat) 0.01%.] ▶? ♀C ▶? \$\$

**PILOCARPINE—OPHTHALMIC** (*Isopto Carpine*) 1 gtt in each affected eye up to four times per day. [Generic/Trade: Soln 1, 2, 4% (15 mL).] ▶Plasma ♀C ▶? \$\$\$

**Glaucoma Agents—Prostaglandin Analogs**

**BIMATOPROST** (*Lumigan, Latisse \* Vistitan*) **Glaucoma** (Lumigan): 1 gtt in each affected eye at bedtime. **Hypotrichosis of the eyelashes** (Latisse): 1 gtt to eyelashes at bedtime. [Generic only: Soln 0.03% 2.5, 5, 7.5 mL. Trade only: Soln 0.01% (Lumigan), 2.5, 5, 7.5 mL. Generic/Trade: Soln 0.03% (Latisse) 3 mL with 70 disposable applicators, 5 mL with 140 disposable applicators.] ▶LK ♀?/?/? ▶? \$\$\$\$

**LATANOPROST** (*Xalatan*) 1 gtt in each affected eye at bedtime. [Generic/Trade: Soln 0.005% (2.5 mL).] ▶LK ♀C ▶? \$

**LATANOPROSTENE BUNOD** (*Vyzulta*) **Chronic open-angle glaucoma or ocular HTN:** 1 gtt in affected eye once daily in the evening. [Trade only: Soln 0.024%, 5 mL.] ▶Minimal absorption ♀?/?/? ▶? \$\$\$\$\$

**TAFLUPROST** (*Zioptan \* Saflutan*) 1 gtt in each affected eye q pm. [Trade only: Soln 0.0015%] ▶L ♀C ▶? \$\$\$\$\$

**TRAVOPROST** (*Travatan Z*) 1 gtt in each affected eye at bedtime. [Trade only: Benzalkonium chloride-free (Travatan Z) 0.004% (2.5, 5 mL).] ▶L ♀C ▶? \$\$\$

**Glaucoma Agents—Sympathomimetics**

**NOTE:** Do not administer while wearing soft contact lenses. Wait 10 min after use before inserting contact lenses. On average, each mL of eye drop soln contains approximately 20 gtt. Reserve ointment formulations for bedtime use due to severe vision blurring. Most eye medications can be administered 1 gtt at a time despite common manufacturer recommendations of 1 to 2 gtt

(cont.)